

Box- and peanut-shaped bulges [★]

I. Statistics

R. Lütticke, R.-J. Dettmar, and M. Pohlen

Astronomisches Institut, Ruhr-Universität Bochum, D-44780 Bochum, Germany
email: luettt@astro.ruhr-uni-bochum.de

Received 11 May 2000; accepted 14 June 2000

Abstract. We present a classification for bulges of a complete sample of ~ 1350 edge-on disk galaxies derived from the RC3 (*Third Reference Catalogue of Bright Galaxies*, de Vaucouleurs et al. 1991). A visual classification of the bulges using the Digitized Sky Survey (DSS) in three types of b/p bulges or as an elliptical type is presented and supported by CCD images. NIR observations reveal that dust extinction does almost not influence the shape of bulges. There is no substantial difference between the shape of bulges in the optical and in the NIR. Our analysis reveals that 45 % of all bulges are box- and peanut-shaped (b/p). The frequency of b/p bulges for all morphological types from S0 to Sd is $> 40\%$. In particular, this is for the first time that such a large frequency of b/p bulges is reported for galaxies as late as Sd. The fraction of the observed b/p bulges is large enough to explain the b/p bulges by bars.

Key words: Galaxies: evolution – Galaxies: spiral – Galaxies: statistics – Galaxies: structure

1. Introduction

Several theories of bulge formation are currently discussed: primordial scenarios (monolithic collapse: Eggen et al. 1962 (ELS); clumpy collapse: Kauffmann et al. 1993, Baugh et al. 1996; inside-out formation: van den Bosch 1998, Kepner 1999), secular evolution scenarios (merger: Wyse et al. 1997 and references therein, dynamical evolution due to gravitational instabilities, such as bars: Combes et al. 1990), and combinations of both (Combes 2000). Therefore statistics of bulges are needed to test the proposed dynamical processes of bulge formation. Statistics of b/p bulges are of particular interest to demonstrate the importance of these structures for disk galaxies and to point out the relevant evolution scenarios of bulges in general. From the frequency of b/p bulges the likeliness of their formation process can be estimated.

External cylindrically symmetric torques (May et al. 1985) or mergers of two disk galaxies (Binney & Petrou 1985, Rowley 1988) as origins of b/p bulges require very special conditions (Bureau 1998). Therefore such evolutionary scenarios can explain only a very low frequency of b/p bulges. Accretion of satellite galaxies is a formation process of b/p bulges (Binney & Petrou 1985, Whitmore & Bell 1988), which could produce a higher frequency of b/p bulges. However, an oblique impact angle of the satellite is needed for the formation of a b/p bulge and a massive accretion event would disrupt the stellar disk (Barnes 1992, Hernquist 1993). The origin of b/p bulges by bars was first noticed by Combes & Sanders (1981). Combes et al. (1990) and Raha et al. (1991) substantially revived this idea. Comparisons between the frequency of barred galaxies and galaxies with a b/p bulge can test the probability of evolutionary scenarios of b/p bulges based on bars.

The estimated frequency of galaxies with a b/p bulge has steadily increased in former statistics from 1.2 % (Jarvis 1986), over 13 % (de Souza & dos Anjos 1987, hereafter SA87) and 20 % (Shaw 1987), up to 45 % (Dettmar & Barteldrees 1988). This rise results from differences in sample selection, sample size, detection method, and criteria to identify b/p bulges. However, the later three statistical studies have shown that b/p bulges are not really as peculiar as they were supposed to be in the past. Therefore very common processes are required to explain their high frequency. No complete statistics and list of galaxies with b/p bulges, based on observations, have been published in the last ten years. However, the knowledge about galactic evolution would greatly benefit from a reliable value for the frequency of b/p bulges by the determination of the most likely scenario for the formation of b/p bulges and subsequent conclusions on the evolution of bulges in general. A complete list of b/p bulges is a base for a detailed analysis of these structures.

In *Sect. 2* we present our investigated sample of edge-on disk galaxies, and our classification of the bulge shape is explained in *Sect. 3*. The results of our statistics and of additional follow-up CCD observations in the optical

[★] Partly based on observations collected at ESO/La Silla (Chile), DSAZ/Calar Alto (Spain), and Lowell Observatory/Flagstaff (AZ/USA).

and in the NIR are given in *Sect. 4*. *Sect. 5* compares our findings to former statistics. In *Sect. 6* we discuss our results and finally, in *Sect. 7* we give the conclusions from our statistical analysis of the sample.

2. Sample selection

2.1. Former samples

The first search for b/p bulges was conducted by Jarvis (1986). By visual inspection of all fields of the ESO/SERC *J* sky survey he found therein 30 b/p bulges and listed additionally 11 galaxies with b/p bulges from the literature. Together with a control sample he estimated that about 1.2% of all disk galaxies possess b/p bulges.

Shaw (1987) inspected visually the ESO/SERC *J* and Palomar Observatory/National Geographical Survey (POSS) *B*- or *R*-band scanned survey plates. He derived a frequency of b/p bulges of $20(\pm 4)\%$ as a lower limit detecting 23 b/p bulges in a sample of 117 disk galaxies. The sample was mainly selected from the RC2 (*Second Reference Catalogue of Bright Galaxies*, de Vaucouleurs et al. 1976) and included disk galaxies with a diameter $\log D_{25} \geq 1.55$ ($D_{25} \sim 3.5'$) at the 25 (*B*) mag arcsec $^{-2}$ isophotal level and a “sufficiently” edge-on aspect.

Until today the largest investigated sample of 555 galaxies was presented by SA87. Their list was extracted from several catalogues and contained all disk galaxies with restricted total *B* magnitude ($B_T < 13.2$ mag) and axis ratio ($b/a < 0.5$). The galaxies were inspected on film copies of the ESO Quick Blue survey or the POSS prints by means of a microscope. A “substantial number” of candidates were scanned using the PDS microdensitometer at ESO and checked with an image processing system (IHAP). In this way SA87 found 74 b/p galaxies as a lower limit. This is 13% of their total sample.

A much larger frequency of b/p bulges was proposed after the investigation of 73 galaxies obtained with CCD surface photometry (Dettmar & Barteldrees 1988, Dettmar 1989). That sample gave a frequency of $45(\pm 8)\%$ and Dettmar (1996) concluded from this sample a lower limit of the total frequency of b/p bulges of 35%. However, only 15 objects with b/p bulge from this sample were listed by name in a paper of Shaw et al. (1990) together with other known b/p bulges from the literature.

2.2. New sample selection

The former statistics in mind a new one should contain a large number of galaxies, be complete, and based on a detection method for the b/p bulges with reasonable accuracy. In the new statistics presented here the completeness with regard to the size of a galaxy is reached by the selection of all disk galaxies ($-3.5 < T < 9.5$) out of an electronic version of the RC3 with diameters larger than $2'$ ($\log D_{25} \geq 1.3$). For disk galaxies without known *T* parameter it is determined separately (see notes to Table 6).

Extracting the sample from one catalogue is preferable to the method of SA87 since the parameters differ among the catalogues and an exact selection criterion is impossible. Similar to Shaw (1987) the selected sample is diameter limited, but compared to his sample the limit is nearly 2 times smaller. Additionally, the use of the newer catalogue (RC3 instead of RC2) leads to a much larger number of investigated galaxies. Restricting the sample in diameter and not in B_T magnitude is preferred because the RC3 attempts to be complete for galaxies with an apparent diameter larger than $1'$ at the D_{25} isophotal level and in addition, the B_T magnitude is not even listed for all galaxies larger than $2'$. Therefore a magnitude limit would not result in a complete sample. Furthermore, a magnitude limit would prefer the selection of early type galaxies.

Since b/p bulges can only be observed at inclinations down to $i \sim 75^\circ$ (Shaw et al. 1990), in a first step all face-on galaxies are excluded using the axis ratio according to SA87. S0 galaxies ($-3.5 < T \leq -0.5$) with $\log R_{25} \geq 0.30$ and other disk galaxies with $\log R_{25} \geq 0.35$ are included ($R_{25} = \frac{b}{a}$). These different limits are used since the transformation of the axis ratio into the inclination angle depends on the morphological type (Bottinelli et al. 1983, Guthrie 1992). With different formulae (the simplest is $\cos i = \frac{b}{a}$; see references in Guthrie 1992) this restriction results for disk galaxies in a limit for the inclination angle i between 60° and 70° degree. This limit ensures most likely a detection of all observable b/p bulges. The final sample meeting these selection criteria contains 1343 galaxies.

The preferable method to classify a b/p bulge is the use of CCD images, but for the whole sample the amount of observing time would be unreasonably large. Therefore all galaxies are inspected using the Digitized Sky Survey (DSS)¹. This survey is complete over the whole sky and an investigation of the images of the galaxies with data analysis software is possible. The DSS is based on photographic surveys of the northern POSS *E* plates (*R*-band, $m_{\text{lim}} = 20.0$ mag), the southern SERC *J* plates (equivalent to *B*-band, $m_{\text{lim}} = 23.0$ mag), and the southern Galactic plane SERC *V* plates (*V*-band, $m_{\text{lim}} = 14.0$ mag) (McLean 1999) and has a scale of $1.7'' \text{pixel}^{-1}$. Galaxies which are saturated in their central regions are checked (if possible) with the ESO Lauberts-Valentijn Archive (Lauberts & Valentijn 1989) kindly made available by ESO². While the images within the ESO Lauberts-Valentijn Archive are not saturated and have a better scale ($1.35'' \text{pixel}^{-1}$), they have a lower signal-to-noise ratio compared to the images of the DSS.

3. Classification of box/peanut bulges

Shaw (1993a) and Lütticke (1996) derive objective parameters to characterize quantitatively b/p structures by

¹ <http://arch-http.hq.eso.org/cgi-bin/dss>

² <http://archive.eso.org/wdb/wdb/eso/esolv/form>

the measurement of the excess luminosity and the total or maximal fraction of the b/p distortion (depending on the radial distance from the center) in relation to the observed bulge luminosity. However, the application of such a classification method is unpractical for a large sample of galaxies and strongly depends on the modelled luminosity distribution for the disk and the elliptical part of the bulge.

The a_4 isophote shape parameter is used by Combes et al. (1990), Shaw (1993a), Lütticke (1996), and Merrifield & Kuijken (1999) to quantify the degree of boxiness of bulges by measuring the deviations from perfect ellipses (e.g. Bender & Möllenhoff 1987). However, it is problematic that the determination of the extreme value of a_4 depends on the fitted region of the bulge. Fitting only the inner parts of bulges results in some galaxies in the undetection of the boxiness of the bulge which is most prominent in the outer parts. For instance the clear boxy bulge of NGC 1055 (Shaw 1993b, Lütticke et al. 2000b, hereafter Paper III) is undetected by Merrifield & Kuijken (1999) using the a_4 parameter. An additional disadvantage of the determination of this parameter is the influence of the masked stars in the foreground, dust, bars, and the extreme nature of b/p isophotal distortions for the ellipse fitting. Therefore the region of the galaxy which is used for the fitting is important. Shaw (1993a) uses the whole galaxy, Lütticke (1996) the disk subtracted galaxy, and Merrifield & Kuijken (1999) mask out a wedge-shaped region of each image within 12 degrees of the disk major axis and only fit “on the side of the galaxy where the disk projects behind the bulge”.

These disadvantages lead to the fact that the a_4 parameter is not suitable for uniform classification of a large sample of bulges. Additionally, the resolution of the DSS images is too low to determine the a_4 parameter for the whole sample. Therefore the bulges of the investigated sample are classified by their degree of b/p shape derived by visual inspection from contour plots of the galaxy images.

Reshetnikov & Combes (1998) use similar arguments to choose a “straightforward procedure of eyeball estimation” with isophotal maps from the DSS instead of objective criteria for the detection and classification of warped disks.

The bulges are divided in three types of b/p bulges (**1** - **3**), elliptical bulges, and unclassifiable bulges.

- 1** : peanut-shaped bulge
- 2** : box-shaped bulge
- 3** : bulge is close to box-shaped, not elliptical
- 4** : elliptical bulge
- 5** : unclassifiable bulge

Type **1** bulges are described by a depression along the minor axis on both sides of the main axis. In this way the bulge looks like a peanut (Fig. 1, top). The depth

of the depression can be used as a characteristic parameter for these peanut-shaped bulges (Lütticke 1999). The box-shaped bulges (type **2**) are defined by isophotes parallel to the major axis. Therefore the bulge appears like a box (Fig. 1, second image). Some bulges of this class have a very prominent box form (Fig. 3, top). They could be type **1**, but due to low resolution of the images it is not possible to see the eventual depression along the minor axis. These bulges are called **2+**. The same class is used for bulges, which are on one side boxy and on the other peanut-shaped. Type **3** (Fig. 1, third image) is less well defined. These bulges possess a general flattening of the isophotes parallel to the major axis which is less pronounced as the flattening of type **2**. However, this flattening differentiates these bulges from the purely elliptical bulges of type **4** (Fig. 1, bottom). The limits between the classes are not sharp: obvious between **1** and **2**, clear between **2** and **3**, but sometimes indistinct between **3** and **4**. The classification of bulges can therefore in some cases be ambiguous, but a check of the whole sample (two independent classifications by the author and one of the co-authors) shows that more than 90% of the bulges are well defined by one of the different bulge types.

Several reasons are differentiated why bulges are not classifiable:

- 5.1** : Inclination is too far away from edge-on.
- 5.2** : Stars in the foreground projected onto the bulge.
- 5.3** : Galaxy is strongly perturbed.
- 5.4** : Dust conceals the shape of the bulge.
- 5.5** : Bulge is too small.
- 5.6** : Signal-to-noise ratio of the image is too low.

For galaxies which meet more than one of the above reasons only the most important one is listed (Table 6 and 7). The classification for all galaxies in the sample with some minor changes after inspection of CCD images (Sect. 4.2) are given in Table 6.

4. Results

4.1. Statistics

About 3/4 of the Sdm and Sm galaxies are not at all classifiable and the rest is highly uncertain, because the bulges of these galaxies — if they have any — are very small and faint and the influence of dust is very strong. Other structures like bars or star forming regions are more dominant. Therefore the selected sample is reduced to disk galaxies ranging from S0-Sd ($-3.5 < T < 7.5$) resulting in 1224 galaxies which will be investigated in the following.

40.0% of these bulges are unclassifiable, almost 3/4 of them due to an inclination which is too far from edge-on. This is an expected effect since the selection criterion of our sample for the inclination is a lower limit (see above). Only a few galaxies with a classifiable bulge in the sample have an axis ratio which is near to the sample




Fig. 1. Examples for the different bulge types from the DSS. Isophotes are logarithmically equidistant.

limit of $\log R_{25} = 0.30$ (S0) and $\log R_{25} = 0.35$ (Sa - Sd), respectively. This shows that the limits are well chosen. There are only two peculiar edge-on galaxies known (with $\log D_{25} > 2'$) which have classifiable bulges and are not in our sample due to their low axis ratio (IC 2560 and NGC 7123). Therefore our list of galaxies with b/p bulges and a diameter larger than $2'$ is almost complete and the number of non-included b/p bulges is small.

Stars in the foreground (type 5.2) are responsible for 13 % of unclassifiable bulges. These stars can influence the projected light from the galaxy in a way that the shape of the bulge cannot be identified. Dust lanes (5.4) prevent classifications in the same way, but only for a few galaxies, in which the dust lane has a large extension in comparison with the bulge size (see also Sect. 4.3). Their fraction decreases from late types and is equal to zero for galaxies earlier than Sbc. Scd and Sd galaxies have too small bulges (5.5) for classification, if their diameter is around the limit of $2'$ and their inclination is near to 90° . At this orientation the measured diameter (D_{25}) and therefore the ratio of disk to bulge length is maximal due to optical depth effects (Xilouris et al. 1999). The result is that galaxies of the same morphological type and diameter (D_{25}) have bulges of different sizes which depend on the inclination angle. However, the fraction of galaxies in the sample having such small bulges is negligible (Table 1). In a few cases galaxies are in the southern Galactic plane (5.6), where the DSS has a higher surface brightness limit due to the use of the SERC V plates (see above). Therefore the images of the galaxies in this region have frequently a very small signal-to-noise ratio and the bulges appear to be too faint for classification. Perturbations (5.3) by interactions are only in a very few cases the reason for bulges being not classifiable.

Table 1. Reasons for the unclassifiability of bulges

bulge type	5.1	5.2	5.3	5.4	5.5	5.6
frequency	74 %	13 %	2 %	4 %	4 %	3 %

Table 2. Classifiable bulges binned by morphological type

	1	2 ¹	3	4	Σ
S0/S0a	5	21	28	78	132
Sa/Sab	6	13	23	52	94
Sb/Sbc	14	44	65	136	259
Sc/Scd	5	28	53	100	186
Sd	0	9	16	38	63
Σ	30	115	185	404	734
%	4.1	15.7	25.2	55.0	100

¹: 10.4 % of all type **2** are **2+**.

734 galaxies of our sample are classifiable (Table 2). From these galaxies we get a frequency of $45.0(\pm 4.5)\%$ b/p bulges (type **1** + **2** + **3**) (Table 2). The distribution of all galaxies with b/p bulges binned by morphological type shows a weak maximum at Sb/Sbc galaxies (Fig. 2). The smallest fraction of b/p bulges is observed for early and late type disk galaxies. The fractions range from 40 to 48 % in the maximum. From number statistics the errors are 4 - 7 % for the frequencies of b/p bulges in each bin. Within these errors there is no dependence of the morphological type discernable. Regarding only galaxies with b/p type **1** and **2** the distribution is nearly the same. These results are in some aspects in contrast to the former investigations (see Sect. 5).




Fig. 2. Frequency of b/p bulges binned by morphological type derived from Table 2: Dotted lines: type **1**. Dashed lines: types **1** + **2**. Solid lines: all b/p types (**1** - **3**).

4.2. CCD observations

The optical CCD images were obtained in several observing runs between 1985 and 1998 at Lowell Observatory (1.06m), Calar Alto (1.23m), and ESO/ La Silla (0.9m, 1.54m, 2.2m, and NTT). Standard reduction techniques for bias subtraction and flatfielding were applied. Individual short exposure frames of the galaxy were combined. The data are partly published in Barteldrees & Dettmar (1994), and Pohlen et al. (2000). Further data will be reported together with follow-up observations of b/p bulges in a separate paper (Lütticke et al. in prep.). In total we have observed 74 galaxies of our investigated RC3 sample up to now.

Only small differences ($< 10\%$) between the classification of bulges comparing DSS (RC3 sample) and CCD images (Fig. 3 or Fig. 1 [top] and Fig. 4 [top]) are detected. Three bulges turned out to be type **4** rather than type **3** and two bulges are type **3** rather than type **4**. This could be explained by the higher resolution of the CCD images

and the unsharp nature of the border between these two classes. However, the total frequency of b/p bulges is not changed. Bulges of type **2+** in the DSS images can be classified more accurately by the high resolution of the CCD images. If there is any depression along the minor axis (e.g. NGC 1886, Fig. 3), the bulge type is changed to **1**. If not, the bulge can be classified as (type **2**). Bulges which have only a depression along the minor axis on one side are still classified as type **2+**.




Fig. 3. Inspecting the DSS image (top) it is hard to decide, if there are depressions along the minor axis of NGC 1886 (first classified as bulge type **2+**) on both sides of the galaxy plane. The contour plots of the CCD images (NGC 1886: ESO/2.2m, 30min in r) reveal that NGC 1886 has a peanut bulge (bulge type **1**).

The comparison of the CCD with the DSS images shows that neither the low resolution, nor the saturated bright parts of some galaxies, nor the low signal-to-noise ratio, nor unresolved structures (small stars in the foreground) strongly influence the classification. Therefore the results of the statistics, which are derived from the RC3 sample and inspected with the DSS, are strongly supported by the CCD observations. Furthermore, the 33

galaxies observed in different optical filters reveal the same shape of the bulge. Therefore the classification of bulges at optical wavelengths is independent of the filter.

47 bulges of galaxies not included in the investigated RC3 sample (45 have $D_{25} < 2'$, additionally, one galaxy has $\log R_{25} < 0.35$, and one S0 galaxy has $\log R_{25} < 0.30$) are classified on CCD images of our observing runs (Table 7).

4.3. NIR observations

The essential advantage of investigations in the NIR is the small extinction by dust at these wavelengths (Knapen et al. 1991). Therefore the possible influence of the dust lane near the galactic plane in edge-on galaxies on the bulge shape is largely reduced by NIR observations. Our NIR sample of galaxies with classifiable bulges consists of 60 galaxies (Lütticke et al. 2000a, hereafter Paper II). It reveals that 75% of the bulges have the same bulge type in the optical as well as in the NIR (Table 3). 21% of the bulges are classified in the NIR to the next lower class, these bulges are less box-shaped. The change of the bulge type is for two bulges in the opposite way, one of them from a boxy to a peanut bulge likely due to the low resolution and saturation of the DSS image of this galaxy (NGC 5166).

Table 3. Differences in the classification of bulges in the optical and NIR

opt. bulge type	4	4	3	3	3	2	2	2	1	1
NIR bulge type	4	3	4	3	2	3	2	1	2	1
no. of galaxies	12	1	7	9	0	5	12	1	0	9

All peanut bulges prominent in the optical have the same shape in the NIR (e.g. ESO 443-42, Fig. 4). A detailed inspection of the galaxies having less boxy bulges in the NIR shows that dust is a possible explanation only for up to one third of these bulges. Differences in classification can be explained mainly by general uncertainties in the classification, the low resolution in the DSS — there is no difference between classifications derived from optical CCD and NIR images —, and low quality of some NIR images (bad seeing). The bulges whose classes are changed can frequently be marked as intermediate types (type **2-3** and **3-4**), respectively. Additionally, seeing conditions around $3''$ weaken the boxy structure and over 50% of the bulges whose class changes were observed under such conditions.

Images of galaxies observed in all three NIR filters J , H , and K reveal no difference in the bulge shape. Therefore the bulge classification seems to be same in a large region of wavelengths from the optical (350 nm) to the NIR (2.2 μm).




Fig. 4. The peanut bulge of ESO 443-42 is prominent in optical CCD images as well as in the NIR. Top: 0.9m/ESO, 10min in R . Bottom: 2.2m/ESO, 23min in K' .

5. Comparison with former statistics

The low frequency of b/p bulges (1.2%) in the sample of Jarvis (1986) seems to result from the limitation to objects of very small diameters (Shaw 1987). Such a restriction, coupled with the used visual detection method on the plates of the ESO/SERC J sky survey, results in a non-detection of many b/p bulges and in the erroneously small frequency of b/p bulges. However, his division in box (**B**) and peanut-shaped (**P**) bulges is in agreement with our classification of b/p bulges. Considering only galaxies included in both samples, 3/4 of the galaxies of type **P** correspond to our bulge type **1** and type **B** to bulge type **2**, respectively. All galaxies with b/p bulges listed by Jarvis (1986) are also classified here as b/p bulges. However, his statistics are based on a too small number of galaxies being representative. Additionally, his sample is biased with galaxies of known b/p bulge characteristics from the literature, therefore the conclusion with regard to a dependence of b/p bulges on the Hubble type is not meaningful.

Shaw (1987) uses a similar definition for his selection criterion of b/p bulges (depression along the minor axis or a general flattening of the isophotes) as we do, but without classification. The low frequency of 20 %, he states, can be explained by the fact that galaxies with b/p bulges of type **3** are almost completely neglected. Counting only type **1** and **2** in our investigated sample results also in a frequency of 20 % (Table 2). However, some prominent b/p bulges are missed by excluding peculiar disk galaxies (e.g. NGC 3628), galaxies which are far from edge-on (e.g. NGC 7582: Quillen et al. 1997), or by simply ignoring others (e.g. NGC 2424: Pohlen et al. 2000). On the other side there are two galaxies (NGC 1596 and NGC 4958) in his list of b/p bulges which have obviously an elliptical bulge. The sample also shows a maximum in the distribution of galaxies with b/p bulges concerning the morphological type of these galaxies at Sb/Sbc galaxies (Table 5). However, the frequencies in the individual bins are much smaller reflecting the total low frequency of b/p bulges. Additionally, the lack of Sd galaxies and the small frequency of Sc/Scd galaxies with b/p bulges are remarkable. That is in contrast to the rather high percentage of late type galaxies with b/p bulges which we have found in our study. The low frequency of Sc/Scd galaxies and the lack of Sd galaxies with b/p bulges are likely a result of the faint nature of bulges and therefore also of their b/p structures. Our method of detection using the DSS and new data analysis systems could explain the differences to Shaw (1987) in the bins of the late types, although both investigations are based on the same photographic material.

SA87 define three classes of galaxies marked by the degree of separation between disk and bulge, and by the form of the bulge. Additional to the mixture of these two criteria, it must be mentioned that their class III is defined as “cigar shaped” bulge. Therefore this class has nothing in common with b/p bulges. This reveals their prototype for class III (NGC 1380) and the high fraction of elliptical bulges (68 %) in the class III sample (Table 4). Therefore the title of their list “box-shaped galaxies” is misleading. However, it should be clear by their definition of the classes that they do not present a list and classification of b/p bulges. This was and is still misunderstood in the literature (e.g. Bureau & Freeman 1999). Therefore a strong correlation between their types and our new introduced bulge types of the galaxies in both samples cannot be expected and is indeed not present (Table 4). However, their type I and I-II (“clear rectangular or peanut shape”) should be classified also in our system as b/p bulge, but four galaxies ($\sim 25\%$) out of these types have clear elliptical bulges (e.g. NGC 6504, Fig. 1, bottom). They also missed, likely due to the bad resolution of the images, some bright galaxies with prominent b/p bulges (e.g. NGC 3079: Shaw et al. 1993, Veilleux et al. 1999; NGC 7582: Quillen et al. 1997). Therefore their list is far from being complete and the frequency of b/p galax-

ies derived by SA87 is not comparable to the frequency of b/p bulges derived in our investigation. The statistics with regard to the Hubble type are not comparable due to the different definition of b/p bulges. In this way it is not surprising that the maximum in the distribution of b/p galaxies is reached for S0 galaxies and clearly lower values for intermediate types (Sa/Sb) in SA87 (Table 5) are not detectable in our distribution of b/p bulges.

Table 4. Our classification compared to SA87’s classification

		bulge type			
		1	2	3	4
galaxy class from SA87	I	6	2	4	4
	II	4	9	8	12
	III	0	2	4	13

Listed are the numbers of galaxies.

Our derived frequency of b/p bulges is in excellent agreement with the result of Dettmar & Barteldrees (1988) and Dettmar (1989) who find a frequency of $45(\pm 8)\%$. Their detection method by use of CCD images seems to prevent a misclassification. All five galaxies in their sample of b/p bulges (published in Shaw et al. 1990) and our sample have bulge types **1** or **2**. Dettmar’s (1989, 1996) statistics concerning the Hubble type reveal also a maximum in the distribution of galaxies with b/p bulges at Sb/Sbc galaxies. However, the differences between the individual bins are much larger than in our statistics (Table 5). This can be explained by statistical errors due to the small sample size of 73 galaxies. Dettmar (1989, 1996) has also not found any Sd galaxy with b/p bulge in contrast to the frequency of 40 % Sd galaxies with b/p bulges in our study. However, the lack of Sd galaxies is likely a result of the small number of investigated Sd galaxies in his sample. Additionally, the already mentioned faint nature of bulges of late type galaxies and the uncertainties in the classification of bulges increasing to later types (see above) lead to the fact that the boxiness of late type bulges could be missed. Even in small samples, where the possibility of comparison between the different forms of Sd bulges is not given, the detection of b/p bulges could be influenced. No detection of a type **1** b/p bulge in galaxies later than Sc in our sample could have similar reasons, if bulge type **1** in late type galaxies exists at all.

Galaxies in previously studied samples, which do not fulfill the selection criterion of our investigated sample due to morphological misclassifications of the Hubble type or too small diameters or axis ratios, are listed in Table 7. However, most of these galaxies have $D_{25} < 2'$ and therefore their classification with the DSS and ESO Lauberts-Valentijn Archive is uncertain.

Table 5. Comparison of different statistics: frequencies of b/p bulges according to their morphological type

	S0/S0a	Sa/Sab	Sb/Sbc	Sc/Scd	Sd
SA87	33%	13%	15%	3%	— ¹
Shaw ²	25%	33%	36%	8%	0%
Dettmar ³	46%	29%	61%	42%	0%
this study	41%	45%	48%	46%	40%

¹: no Sd galaxy in sample; ²: frequencies derived from Table 2 of Shaw 1987; ³: frequencies derived from Figure 1 of Dettmar 1989)

6. Discussion

The new classification of bulges is very similar to the types used by Jarvis (1986) and Shaw (1987). The frequency of 45% b/p bulges is consistent with Dettmar & Barteldrees (1988), Dettmar (1989) and the frequency of 20% prominent b/p bulges with Shaw (1987). However, the derived frequency is now based on a much larger sample of 734 galaxies. Furthermore, for the first time a large fraction of b/p bulges in galaxies as late as Sd is found, and some previously unknown peanut bulges (type 1) are listed (Table 6 and 7).

The large fraction of b/p bulges (45%) shows that such bulges are not that peculiar but rather quite normal. Therefore very common processes are required to explain the origin of b/p bulges.

In order to check possible formation processes relating b/p bulges with bars, we have compared the frequency distributions of galaxies with b/p bulges and barred galaxies. The frequency distribution of barred galaxies is derived from RC3 (only face-on galaxies) and contains 8587 galaxies. (Fig. 5). Both distributions binned by morphological type show the same general dependence. The maximum for barred galaxies is also at Sb/Sbc and the minimum at S0/S0a galaxies. This minimum is more distinct for barred galaxies as for galaxies with b/p bulge. However, for all galaxies the fraction of barred galaxies is 55% (nearly 2/3 of them are strongly barred [SB] and the rest weakly barred [SAB = SX]) in relation to 45% of galaxies with b/p bulges. This high percentage of barred galaxies is also recently confirmed by Knapen et al. (2000). Furthermore, it is remarkable that the percentage of barred Sd galaxies is relatively high compared to the minimal frequency of galaxies with b/p bulges for Sd galaxies. However, it should be mentioned that our error of this bin is the largest due to low number statistics (Table 2). Other statistics of barred galaxies taken from older catalogues (Sellwood & Wilkinson 1993) show the same general dependences, although the percentages vary mainly due to different fractions of weakly barred galaxies in the catalogues. However, former samples contain a much smaller number of galaxies (Sellwood & Wilkinson 1993). Unfortunately, the galaxies are not all classified as SB, SX, or

SA (unbarred galaxy), but there is also a group of galaxies without any classification ($S = 28\%$). These galaxies can be likely interpreted as unbarred galaxies (as done in Fig. 5), because in former times unbarred galaxies were simply called “S”, but they add uncertainties to the quantitative conclusion.




Fig. 5. Frequency of barred galaxies binned by morphological type. All face-on galaxies of the RC3 are included, i.e., all galaxies with a diameter (D_{25}) larger than $1'$ and an axis ratio smaller than $1/3$ ($\log R_{25} < 0.477$). Solid lines: strongly barred (SB) plus weakly barred (SX) galaxies. Dashed lines: only SB galaxies.

Bars as origin for b/p bulges (Combes et al. 1990, Raha et al. 1991, Pfenniger & Friedli 1991) are supported by the similarity in frequency distributions of galaxies with b/p bulges and barred galaxies. The fraction of b/p bulges is large enough to explain the b/p bulges by bars. The higher frequency of barred galaxies can easily be explained by the aspect angle of bars. End-on bars result for edge-on galaxies in elliptical shaped bulges, bars with intermediate aspect angles in boxy bulges, and edge-on bars in peanut bulges (Combes et al. 1990, Pfenniger & Friedli 1991, Paper II). Therefore our statistical results are consistent with recent studies stating a strong correlation of bars and b/p bulges. Bureau & Freeman (1999) and Merrifield & Kuijken (1999) find in observations of gas kinematics the characteristic “figure-of-eight” rotation curve, which is a strong signature for the presence of a bar, in many galaxies with b/p bulges. Direct kinematic evidence for streaming motions of a bar in two galaxies with a b/p bulge are reported by Veilleux et al. (1999). Kinematical bar diagnostics in edge-on spiral galaxies using simulations of families of periodic orbits and hydrodynamical simulations confirm the connection between bars and b/p bulges (Bureau & Athanassoula 1999, Athanassoula & Bureau 1999). Additionally, we find in our NIR study (Paper II) a strong correlation of bar signatures with b/p bulges. However, by finding a few bulges with a very complex morphol-

ogy, we do not exclude that some b/p bulges result from a recent merger event (Dettmar & Lütticke 1999, Paper III).

The optical CCD images verify our classification of bulges and thereby the statistics derived by the DSS images. For the analysis of faint structures and more objective parameters of b/p bulges (e.g. parameter for the depression at the minor axis) CCD images are necessary. The best parameter for an objective classification of bulges would be the minimum of the a_4 parameter determined in the NIR (minimizing the dust influence) and after subtraction of a modelled disk and bar. However, models of high quality need images with a high signal-to-noise ratio. This would result in unreasonably observing time for the whole RC3 sample. Therefore the classification by visual inspection seems to be the best method until CCD data of the whole sky are available (e.g. for the northern sky: Sloan Digital Sky Survey).

Furthermore, it is shown by NIR observations that dust is not an important factor for the classification of b/p bulges. Therefore they are indeed present in disks of late type spirals as pointed out previously by Dettmar & Ferrara (1996). This result is not in contrast to Baggett & MacKenty (1996) who verify b/p structures in the NIR only in four out of six galaxies because their sample contains several misclassifications from older literature lists (section 2.1 and 5). The comparison of optical and NIR images of b/p bulges reveals that there is no difference between the shape of bulges in different wavelengths bands or the differences are smaller than the uncertainties of the classification.

Additionally, as a byproduct a new catalogue of edge-on galaxies with $D_{25} > 2'$ confirmed by visual inspection is formed by our classification of bulges. This method gives a homogeneous sample of highly inclined galaxies compared to samples selected only by axis ratio and morphological type. This catalogue includes all galaxies with a bulge of type 1 – 4 and 5.2 – 5.5 (Table 6).

7. Conclusion

The bulge shape of a large sample of edge-on galaxies was investigated using DSS, CCD images in the optical wavelengths regime and in the NIR. Our main conclusions are:

- We have presented a visual classification of bulges in three b/p types and one elliptical type.
- 45 % of all disk galaxies (S0 - Sd) have a b/p bulge. This fraction is based on a sample of 734 galaxies.
- For the first time a large fraction (40 %) of b/p bulges in Sd galaxies is found.
- Comparisons between the fraction of barred galaxies and galaxies with b/p bulges show that b/p bulges can be explained by evolution scenarios based on bars.
- NIR observations reveal that dust is not responsible for the shape of b/p bulges.

- The shape of bulges is constant over a large region of wavelengths (350 nm - 2.2 μ m).

Acknowledgements. Part of this work was supported by the *Deutsche Forschungsgemeinschaft, DFG*. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. It also uses the Digitized Sky Survey (DSS) based on photographic data obtained using Oschin Schmidt Telescope on Palomar Mountain and The UK Schmidt Telescope and produced at the Space Telescope Science Institute.

References

- Athanassoula E., Bureau M., 1999, ApJ 522, 699
- Baggett S.M., MacKenty J.W., 1996, Box/Peanut Galaxies in the Near-IR. In: Buta R., Crocker D.A., Elmegreen B.G. (eds.), ASP Conference Series Volume 91, Barred Galaxies, p. 224
- Barnes J.E., 1992, ApJ 331, 699
- Barteldrees A., Dettmar R.-J., 1994, A&AS 103, 475
- Baugh C.M., Cole S., Frenk C.S., 1996, MNRAS 283, 1361
- Bender R., Möllenhoff C., 1987, A&A 177, 71
- Binggeli B., Sandage A., Tammann G.A., 1985, AJ 90, 1681, VCC
- Binney J., Petrou M., 1985, MNRAS 214, 449
- Bottinelli L., Gouguenheim L., Paturel G., de Vaucouleurs G., 1983, A&A 118, 4
- Bureau M., 1998, Bars in Edge-On Spiral Galaxies, Ph.D. Thesis, The Australian National University
- Bureau M., Athanassoula E., 1999, ApJ 522, 686
- Bureau M., Freeman K.C., 1999, AJ 118, 126
- Combes F., 2000, Bulge Formation. In: Hammer F., Thuan T.X., Cayatte V., Guiderdoni B., Tran Thanh Van J. (eds.), Building Galaxies: from the Primordial Universe to the Present, Proceedings of Rencontres de Moriond, Ed. Frontiers, p. 413
- Combes F., Sanders R.H., 1981, A&A 96, 164
- Combes F., Debbasch F., Friedli D., Pfenniger D., 1990, A&A 233, 82
- de Souza R.E., dos Anjos S., 1987, A&AS 70, 465 (**SA87**)
- Dettmar R.-J., 1989, Box- and Peanut-Shaped Bulges of Disk Galaxies. In: Corwin H.G., Bottinelli L. (eds.), The World of Galaxies, Springer, p.229
- Dettmar R.-J., 1996, Galactic bulges as an evolutionary phenomenon. In: Blitz L., Teuben P. (eds.), IAU Symp. 169, Unsolved Problems of the Milky Way, Kluwer, p. 335
- Dettmar R.-J., Barteldrees A., 1988, BAAS 20, 1085
- Dettmar R.-J., Ferrara A., 1996, NIR imaging of the box/peanut bulge in NGC 4302. In: Bender R., Davis R. (eds.), IAU Symp. 171, New Light on Galaxy Evolution, Kluwer, p. 362
- Dettmar R.-J., Lütticke R., 1999, Do some bulges result from merging ? In: Gibson B.K., Axelrod T.S., Putman M.E. (eds.), ASP Conference Series Volume 165, The Third Stromlo Symposium: The Galactic Halo, p. 95
- de Vaucouleurs G., de Vaucouleurs A., Corwin Jr. H.G., 1976, Second Reference Catalogue of Bright Galaxies (RC2), University of Texas Press

- de Vaucouleurs G., de Vaucouleurs A., Corwin Jr. H.G., Buta R.J., Fouqué P., 1991, Third Reference Catalogue of Bright Galaxies (RC3), Springer
- Eggen O., Lynden-Bell D., Sandage A., 1962, ApJ 136, 748
- Guthrie B.N.G., 1992, A&AS 93, 255
- Haynes M.P., Herter T., Barton S., Benensohn S., 1990, AJ 99, 1740
- Hernquist L., 1993, ApJ 409, 548
- Jarvis B.J., 1986, AJ 91, 65
- Kauffmann G., White S.D.M., Guiderdoni B., 1993, MNRAS 264, 201
- Kepner J.V., 1999, ApJ 520, 59
- Knapen J.H., Hes R., Beckman J.E., Peletier R.F., 1991, A&A 241, 42
- Knapen J.H., Shlosman I., Peletier R.F., 2000, ApJ in press
- Kraan-Korteweg R.C., 1982, A&AS 47, 505
- Lauberts A., 1982, The ESO/Uppsala Survey of the ESO-B Atlas, ESO
- Lauberts A., Valentijn E.A., 1989, The Surface Photometry Catalogue of the ESO-Uppsala Galaxies, ESO
- Lütticke R., 1996, Flächenphotometrische Untersuchungen zu Eigenschaften von "box/peanut"-bulges in "edge-on" Scheibengalaxien, Diploma Thesis, Ruhr-Universität Bochum
- Lütticke R., 1999, Box- and Peanut-Shaped Bulges, Ph.D. Thesis, Ruhr-Universität Bochum
- Lütticke R., Dettmar R.-J., Pohlen M., 2000a, A&A in prep. (Paper II)
- Lütticke R., Dettmar R.-J., Pohlen M., 2000b, A&A in prep. (Paper III)
- May A., van Albada T.S., Norman C.A., 1985, MNRAS 214, 131
- McLean B., 1999, <http://www-gsss.stsci.edu/dss/surveys.html>
- Merrifield M., Kuijken K., 1999, A&A 345, L47
- Pfenniger D., Friedli D., 1991, A&A 252, 75
- Pohlen M., Dettmar R.-J., Lütticke R., Schwarzkopf U., 2000, A&AS 144, 405
- Quillen A.C., Kuchinski L.E., Frogel J.A., Depoy D.L., 1997, ApJ 481, 179
- Raha N., Sellwood J.A., James R.A., Kahn F.D., 1991, Nature 352, 411
- Reshetnikov V., Combes F., 1998, A&A 337, 9
- Rowley G., 1988, ApJ 331, 124
- Sellwood J.A., Wilkinson A., 1993, Reports on Progress in Physics 56, 173
- Shaw M.A., 1987, MNRAS 229, 691
- Shaw M.A., 1993a, MNRAS 261, 718
- Shaw M.A., 1993b, A&A 280, 33
- Shaw M.A., Dettmar R.J., Barteldrees A., 1990, A&A 240, 36
- Shaw M.A., Wilkinson A., Carter D., 1993, A&A 268, 511
- Skiff B., 1999, priv. communication
- van den Bosch F.C., 1998, ApJ 507, 601
- Veilleux S., Bland-Hawthorn J., Cecil G., 1999, AJ 118, 2108
- Whitmore B.C., Bell M., 1988, ApJ 324, 741
- Wyse R.F.G., Gilmore G., Franx M., 1997, ARA&A 35, 637
- Xilouris E.M., Byun Y.I., Kylafis N.D., Paleologou E.V., Pampastorakis J., 1999, A&A 344, 868
- Yasuda N., Okamura S., Fukugita M., 1995, ApJS 96, 359
- Yasuda N., Fukugita M., Okamura S., 1997, ApJS 108, 417

Table 6. List of galaxies in the RC3-sample with bulge classification

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
IC 5376	00 01	+34 31	4			2.0	1.30	0.76
NGC 7814	00 03	+16 08	4 ^f			2.0	1.74	0.38
NGC 7817	00 03	+20 45	5.1			4.0	1.55	0.58
ESO 293- 34	00 06	−41 29	5.4			6.1	1.50	0.51
NGC 7	00 08	−29 54	3			4.5	1.35	0.61
NGC 13	00 08	+33 26	4			2.3	1.39	0.62
NGC 24	00 09	−24 57	3			5.0	1.76	0.63
ESO 241- 21	00 10	−46 25	3			2.8	1.31	0.63
NGC 52	00 14	+18 34	2			2 ¹	1.33	0.70
NGC 59	00 15	−21 26	5.1			−2.5	1.42	0.30
ESO 78- 22	00 20	−63 51	2			4.3	1.31	0.77
NGC 100	00 24	+16 29	4			6.0	1.74	0.90
UGC 240	00 25	+06 29	5.1			3.0	1.39	0.36
MCG -1- 2- 11	00 25	−02 17	4			4.2	1.30	0.90
UGC 260	00 27	+11 35	3			6.0	1.48	0.72
NGC 128	00 29	+02 51	1		J P, SA I, SDB	−2.0	1.47	0.52
NGC 134	00 30	−33 14	2			4.0	1.93	0.62
ESO 473- 25	00 31	−26 43	2			4.7	1.41	0.90
ESO 79- 3	00 32	−64 15	3			3.0	1.43	0.81
NGC 148	00 34	−31 47	4			−1.5	1.30	0.40
IC 34	00 35	+09 07	5.1			1.0	1.45	0.44
ESO 540- 3	00 35	−20 07	5.1			2.8	1.31	0.40
NGC 169	00 36	+23 59	4			2.0	1.42	0.60
NGC 172	00 37	−22 35	2			3.8	1.31	0.79
NGC 177	00 37	−22 32	4			1.8	1.35	0.65
UGC 418	00 39	+08 57	4			3.0	1.33	0.94
NGC 216	00 41	−21 02	4			−2.0	1.31	0.47
NGC 217	00 41	−10 01	3			0.0	1.42	0.63
ESO 540- 16	00 42	−18 09	5.2			6.0	1.43	0.95
NGC 224	00 42	+41 16	4			3.0	3.28	0.49
UGC 484	00 46	+32 40	3			3.0	1.38	0.47
UGC 485	00 47	+30 20	5.5			6.0	1.34	1.01
NGC 247	00 47	−20 45	5.1			7.0	2.33	0.49
NGC 253	00 47	−25 17	4			5.0	2.44	0.61
MCG -2- 3- 16	00 47	−09 53	5.4			6.7	1.48	0.82
NGC 259	00 48	−02 46	3			4.0	1.45	0.66
NGC 273	00 50	−06 53	4			−2.0	1.34	0.52
UGC 542	00 53	+29 16	4			3 ⁶	1.31	0.70
NGC 295	00 55	+31 32	5.1			3.0	1.34	0.36
NGC 328	00 56	−52 55	3			1.4	1.43	0.71
MCG -1- 3- 51	00 59	−04 48	4			1.0	1.32	0.54
IC 1608	00 59	−34 19	5.1			−0.7	1.31	0.36
IC 65	01 00	+47 40	4			4.0	1.59	0.53
NGC 352	01 02	−04 14	3		SA II	3.0	1.38	0.42
NGC 360	01 02	−65 36	3			3.7	1.55	0.87
ESO 13- 12	01 07	−80 18	2			−0.3	1.44	0.51
NGC 406	01 07	−69 52	5.1			5.0	1.52	0.42
IC 1627	01 08	−46 05	3			3.0	1.40	0.60
UGC 711	01 08	+01 38	4			6.7	1.56	0.98
UGC 725	01 10	+43 06	3			6.0	1.35	0.62
MCG -1- 4- 8	01 10	−05 49	5.1			5.0	1.35	0.38
UGC 748	01 11	+35 16	5.1			5 ²	1.33	0.52
IC 1657	01 14	−32 39	4			3.7	1.37	0.63
NGC 452	01 16	+31 02	4			2.0	1.40	0.49
NGC 482	01 20	−40 57	2			2.4	1.35	0.62

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
NGC 493	01 22	+00 56	3			6.0	1.53	0.51
NGC 497	01 22	−00 52	5.1			4.0	1.32	0.38
NGC 494	01 22	+33 10	5.1			2.0	1.30	0.41
NGC 517	01 24	+33 25	5.1			−2.0	1.30	0.30
NGC 522	01 24	+09 59	2			3.7	1.43	0.78
NGC 532	01 25	+09 15	2+			2.3	1.39	0.50
NGC 549	01 25	−38 16	2			−0.5	1.30	0.63
UGC 987	01 25	+32 08	4			1.0	1.34	0.52
MCG -1- 4- 44	01 26	−06 04	5.4			5.1	1.46	0.81
NGC 536	01 26	+34 42	5.1			3.0	1.47	0.44
UGC 1033	01 27	+31 33	5.2			6.0	1.47	0.77
IC 1711	01 30	+17 11	4			3.0	1.41	0.73
NGC 585	01 31	−00 55	3 ^f			1.0	1.33	0.62
NGC 582	01 31	+33 28	3			4 ¹	1.35	0.58
NGC 587	01 32	+35 21	5.1			3.0	1.34	0.43
NGC 615	01 35	−07 20	5.1			3.0	1.56	0.40
NGC 634	01 38	+35 21	4			1.0	1.32	0.52
NGC 645	01 40	+05 43	5.1			3.0	1.42	0.35
NGC 2573	01 41	−89 20	5.1			6.3	1.30	0.42
NGC 660	01 43	+13 38	5.3		Sh	1.0	1.92	0.42
IC 1723	01 43	+08 53	3			3.0	1.52	0.63
NGC 669	01 47	+35 33	2 ^f		SA II	2.0	1.50	0.73
NGC 670	01 47	+27 53	5.1			−2.0	1.31	0.33
NGC 672	01 47	+27 26	5.1			6.0	1.86	0.45
NGC 676	01 48	+05 54	3		SA III	0.0	1.60	0.52
NGC 678	01 49	+21 59	2		SA II, S	3.0	1.65	0.75
NGC 684	01 50	+27 38	3		SA II	3.0	1.51	0.75
NGC 697	01 51	+22 21	5.1			4.7	1.65	0.48
ESO 543- 20	01 52	−20 09	5.1			2.6	1.33	0.38
MCG -1- 5- 47	01 52	−03 26	4			5.0	1.46	0.86
UGC 1400	01 56	+36 07	2			3.0	1.40	0.81
ESO 477- 16	01 56	−22 54	4			3.5	1.31	1.00
NGC 755	01 56	−09 03	3			3.4	1.53	0.49
ESO 245- 10	01 56	−43 58	3			3.0	1.38	0.59
NGC 779	01 59	−05 57	4			3.0	1.60	0.53
IC 187	02 01	+26 28	5.1			1.0	1.31	0.45
MCG -2- 6- 19	02 03	−09 39	4			6.8	1.45	1.10
NGC 801	02 03	+38 15	4			5.0	1.50	0.66
NGC 803	02 03	+16 01	5.1			5.0	1.48	0.38
NGC 825	02 08	+06 19	4			1.0	1.35	0.41
NGC 818	02 08	+38 46	5.1			4.5	1.47	0.36
NGC 827	02 08	+07 58	5.1			3 ²	1.35	0.44
ESO 30- 8	02 09	−75 56	3			7.0	1.35	0.74
IC 210	02 09	−09 40	5.4			4.0	1.36	0.56
UGC 1650	02 09	+37 15	5.5			6.0	1.30	1.31
IC 1783	02 10	−32 56	5.1			2.8	1.30	0.40
MCG -2- 6- 35	02 10	−09 42	5.1			5.0	1.39	0.41
ESO 30- 9	02 10	−75 02	4			5.0	1.40	0.61
MCG -1- 6- 77	02 14	−07 21	4			3.0	1.36	0.81
IC 1788	02 15	−31 12	5.1			4.0	1.41	0.36
IC 217	02 16	−11 55	5.4			6.0	1.34	0.69
IC 1789	02 17	+32 23	4			1.0	1.34	0.78
NGC 876	02 17	+14 31	3			5.0	1.32	0.72
UGC 1817	02 21	+14 12	4			7.0	1.39	0.94
UGC 1839	02 22	−00 37	2			7.3	1.31	0.93
NGC 891	02 22	+42 20	2		J B, SA II	3.0	2.13	0.73

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
NGC 908	02 23	-21 14	5.1			5.0	1.78	0.36
UGC 1856	02 24	+31 36	4			7.0	1.37	0.97
UGC 1883	02 25	+11 28	4			7.0	1.34	0.58
NGC 931	02 28	+31 18	4			4.0	1.59	0.67
UGC 1970	02 29	+25 15	4			6.0	1.37	0.93
NGC 955	02 30	-01 06	4		SA II	1.8	1.44	0.59
NGC 958	02 30	-02 56	5.1			4.5	1.46	0.46
NGC 964	02 31	-36 02	4			2.0	1.31	0.58
UGC 1993	02 31	+39 22	2			3.0	1.37	0.60
UGC 1999	02 31	+19 08	4			6.0	1.49	0.83
NGC 973	02 34	+32 30	2+ ^f			3.0	1.57	0.84
MCG -2- 7- 33	02 34	-10 50	5.1			4.0	1.46	0.44
UGC 2082	02 36	+25 25	4			6.0	1.73	0.81
UGC 2092	02 36	+07 18	4			6.0	1.50	1.06
ESO 355- 30	02 37	-32 55	5.1			4.1	1.30	0.38
NGC 1024	02 39	+10 50	5.1			2.0	1.59	0.43
NGC 1012	02 39	+30 09	4			0.0	1.40	0.36
NGC 1003	02 39	+40 52	5.1			6.0	1.74	0.47
NGC 1032	02 39	+01 05	4		SA II	0.0	1.52	0.47
NGC 1035	02 39	-08 08	5.1			5.0	1.35	0.48
NGC 1023	02 40	+39 03	4			-3.0	1.94	0.47
NGC 1055	02 41	+00 26	2 ^d		SA II, S	3.0	1.88	0.45
UGC 2185	02 43	+40 25	4			6.0	1.47	0.50
ESO 154- 10	02 45	-55 44	5.1			1.5	1.41	0.39
NGC 1090	02 46	-00 14	5.1			4.0	1.60	0.36
NGC 1103	02 48	-13 57	4			3.0	1.33	0.59
ESO 299- 18	02 48	-40 33	4			6.0	1.30	0.95
ESO 416- 25	02 48	-31 32	3 ^d			3.0	1.31	0.82
UGC 2280	02 48	+41 46	5.1			4.0	1.31	0.48
NGC 1118	02 49	-12 09	3			-2.0	1.33	0.48
MCG -2- 8- 12	02 50	-08 35	4			5.0	1.34	0.83
IC 1862	02 51	-33 20	4			3.7	1.48	1.01
NGC 1134	02 53	+13 00	5.1			3 ⁶	1.40	0.46
NGC 1145	02 54	-18 38	2			5.0	1.51	0.80
UGC 2424	02 57	+17 30	2			4.0	1.35	0.59
MCG -2- 8- 33	02 57	-10 10	4			0.0	1.43	0.55
ESO 31- 5	02 58	-74 27	5.1			3.4	1.32	0.40
UGC 2441	02 58	+03 51	3			7.0	1.34	0.72
NGC 1165	02 58	-32 05	2+			0.7	1.40	0.42
NGC 1163	03 00	-17 09	4			3.7	1.35	0.85
NGC 1200	03 03	-11 59	5.1			-3.0	1.47	0.30
NGC 1171	03 03	+43 23	5.1			6.0	1.42	0.36
ESO 248- 2	03 05	-45 57	4			7.0	1.48	0.66
NGC 1186	03 05	+42 50	5.1			4.0	1.50	0.42
UGC 2526	03 05	+36 47	4			3.0	1.55	0.68
IC 1898	03 10	-22 24	2			5.5	1.56	0.80
NGC 1247	03 12	-10 28	3			4.0	1.53	0.81
ESO 116- 12	03 13	-57 21	5.4			6.5	1.54	0.51
NGC 1253	03 14	-02 49	5.1			6.0	1.72	0.35
NGC 1250	03 15	+41 21	4			-1.7	1.33	0.40
MCG -2- 9- 19	03 15	-12 01	4			-1.0	1.35	0.57
UGC 2617	03 16	+40 53	5.1			7.0	1.40	0.45
NGC 1184	03 16	+80 47	3		SA II	0.0	1.45	0.67
NGC 1292	03 18	-27 36	5.1			5.0	1.47	0.35
ESO 481- 18	03 18	-25 50	5.1			5.5	1.46	0.46
UGC 2679	03 20	+01 21	3			3.0	1.30	0.59

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
NGC 1301	03 20	−18 42	2			3.5	1.34	0.69
UGC 2709	03 23	+38 40	4			3.0	1.37	0.48
NGC 1325	03 24	−21 32	5.1			4.0	1.67	0.47
NGC 1324	03 25	−05 44	3			3.0	1.32	0.42
ESO 548- 16	03 26	−21 20	5.1			5.0	1.37	0.53
NGC 1332	03 26	−21 20	4		SA III	−3.0	1.67	0.51
NGC 1337	03 28	−08 23	4			6.0	1.76	0.59
NGC 1351A	03 28	−35 10	4			4.0	1.43	0.70
UGC 2752	03 29	+40 49	4			−2.0	1.30	0.62
NGC 1353	03 32	−20 49	5.1			3.0	1.53	0.38
UGC 2765	03 32	+68 22	5.1			0 ⁶	1.60	0.40
NGC 1354	03 32	−15 13	3			0.0	1.35	0.47
IC 1952	03 33	−23 42	5.2			4.0	1.41	0.64
NGC 1366	03 33	−31 11	4		SA III	−2.0	1.32	0.35
ESO 548- 47	03 34	−19 01	2			−1.3	1.39	0.59
NGC 1375	03 35	−35 15	2			−2.0	1.35	0.42
IC 335	03 35	−34 26	4			−2.0	1.41	0.59
NGC 1380	03 36	−34 58	4		SA III	−2.0	1.68	0.32
NGC 1381	03 36	−35 17	2		J B, SA I, SDB	−1.6	1.43	0.56
IC 1970	03 36	−43 57	4			2.6	1.50	0.63
NGC 1386	03 36	−35 59	4			−0.6	1.53	0.42
NGC 1380A	03 36	−34 44	2			−2.0	1.38	0.54
NGC 1401	03 39	−22 43	3			−2.0	1.38	0.57
NGC 1406	03 39	−31 19	2			4.0	1.58	0.70
NGC 1422	03 41	−21 40	3			2.4	1.35	0.61
UGC 2817	03 41	+37 12	5.1			7 ²	1.32	0.45
NGC 1425	03 42	−29 53	5.1			3.0	1.76	0.35
NGC 1421	03 42	−13 29	2			4.0	1.55	0.61
MCG -2-10- 9	03 42	−12 54	4			5.0	1.49	0.82
MCG -3-10- 42	03 44	−14 21	5.1			4.0	1.31	0.47
NGC 1448	03 44	−44 38	4			6.0	1.88	0.65
NGC 1438	03 45	−23 00	5.1			0.0	1.30	0.36
MCG -1-10- 35	03 46	−04 27	4			5.0	1.37	0.85
NGC 1461	03 48	−16 23	4			−2.0	1.48	0.51
IC 2000	03 49	−48 51	5.4			5.5	1.61	0.70
ESO 482- 46	03 49	−26 59	3			5.3	1.53	0.83
ESO 156- 18	03 52	−54 53	5.1			−2.0	1.33	0.30
NGC 1484	03 54	−36 58	3			3.0	1.40	0.64
NGC 1495	03 58	−44 27	4			5.0	1.48	0.74
NGC 1511	03 59	−67 38	4			0.7	1.54	0.46
ESO 483- 6	04 00	−25 10	3			3.2	1.42	0.76
UGC 2920	04 00	+35 00	4			6.0	1.36	0.82
UGC 2936	04 02	+01 57	5.4			7.0	1.40	0.57
NGC 1515	04 04	−54 06	4			4.0	1.72	0.67
NGC 1485	04 05	+70 59	3			3.0	1.33	0.49
UGC 2963	04 07	+03 58	5.1			7.0	1.30	0.51
NGC 1519	04 08	−17 11	2			3.4	1.33	0.59
NGC 1527	04 08	−47 53	4			−2.7	1.57	0.43
ESO 201- 22	04 09	−48 43	4 ^d			5.0	1.40	0.87
NGC 1532	04 12	−32 52	2			2.7	2.10	0.58
UGC 2983	04 12	+02 22	4			3.3	1.30	0.47
UGC 2988	04 13	+25 28	4			3.0	1.45	0.67
IC 2058	04 17	−55 56	4			6.6	1.47	0.88
NGC 1558	04 20	−45 01	5.1			3.7	1.39	0.38
ESO 550- 24	04 21	−21 50	5.1			6.7	1.75	0.42
NGC 1596	04 27	−55 01	4 ^c		SA I, S	−2.0	1.57	0.59

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
UGC 3042	04 28	+70 20	5.1			4.0	1.33	0.45
NGC 1589	04 30	+00 51	2 ^c		SA III	2.2	1.50	0.49
IC 2085	04 31	−54 25	4			−2.0	1.38	0.67
ESO 202- 35	04 32	−49 40	3			3.0	1.42	0.86
NGC 1560	04 32	+71 52	5.2			7.0	1.99	0.76
MCG -2-12- 41	04 33	−11 42	3			2.0	1.50	0.50
NGC 1618	04 36	−03 08	5.1			3.0	1.37	0.46
NGC 1622	04 36	−03 11	4			2.0	1.56	0.70
NGC 1620	04 36	−00 08	2			4.3	1.46	0.46
NGC 1625	04 37	−03 18	5.3			3.0	1.32	0.62
UGC 3110	04 41	+73 40	5.1			6.0	1.33	0.39
NGC 1645	04 44	−05 27	5.1			−1.0	1.36	0.35
UGC 3137	04 46	+76 25	4			4 ⁶	1.55	0.98
NGC 1670	04 49	−02 45	5.1			−2.3	1.32	0.33
MCG -3-13- 16	04 50	−17 16	2			4.0	1.30	0.59
IC 2098	04 50	−05 25	3 ^d			6.0	1.32	0.88
ESO 361- 15	04 51	−33 10	5.2			7.0	1.47	0.83
MCG -2-13- 21	04 55	−10 41	5.2			3.0	1.46	0.64
UGC 3207	04 56	+02 09	3			2.5	1.37	0.51
UGC 3214	04 57	−00 07	2			3.0	1.51	0.69
NGC 1721	04 59	−11 07	5.1			−2.0	1.32	0.31
NGC 1728	04 59	−11 07	2			1.0	1.30	0.46
NGC 1809	05 02	−69 34	5.2			5.0	1.50	0.59
NGC 1752	05 02	−08 14	5.1			4.5	1.42	0.51
MCG -2-13- 38	05 02	−10 21	5.1			1.0	1.31	0.36
MCG -1-13- 50	05 03	−02 56	4			3.0	1.35	0.76
UGC 3235	05 05	+67 01	5.1			6.0	1.36	0.39
NGC 1827	05 10	−36 57	2			6.0	1.48	0.66
MCG -2-14- 3	05 11	−09 23	3			5.0	1.41	0.68
NGC 1853	05 12	−57 23	5.1			6.7	1.30	0.46
ESO 423- 2	05 15	−30 31	2			7.0	1.45	0.55
ESO 362- 11	05 16	−37 06	4			4.0	1.65	0.78
NGC 1892	05 17	−64 57	5.6			6.0	1.46	0.55
NGC 1886	05 21	−23 48	1 ^d		J P	3.5	1.49	0.85
NGC 1888	05 22	−11 29	3		SA II	5.0	1.48	0.57
ESO 33- 22	05 31	−73 44	5.5			7.0	1.31	1.19
IC 2135	05 33	−36 23	4			5.5	1.44	0.69
NGC 1964	05 33	−21 56	5.1			3.0	1.75	0.42
PGC 17482	05 35	−48 46	5.1			−3.0	1.32	0.50
UGC 3326	05 39	+77 18	4 ^f			6.0	1.55	1.26
ESO 554- 29	05 43	−19 17	5.1			6.1	1.31	0.43
UGC 3343	05 45	+72 21	4			3 ⁶	1.32	0.61
UGC 3354	05 47	+56 06	3	4		2.0	1.32	0.55
ESO 555- 2	05 50	−19 43	3			4.4	1.33	0.80
NGC 2106	05 50	−21 34	5.1			−2.0	1.43	0.30
IC 2150	05 51	−38 19	5.1			4.7	1.42	0.51
ESO 120- 16	05 51	−59 02	4			3.0	1.35	0.75
UGC 3365	05 52	+66 47	3			1.0	1.34	0.88
IC 2160	05 55	−76 55	5.1			4.7	1.31	0.40
NGC 2124	05 57	−20 05	4			3.0	1.43	0.49
ESO 555- 22	06 01	−21 44	5.2			4.0	1.37	0.53
MCG -2-16- 2	06 04	−12 37	5.2			3.0	1.40	0.73
ESO 488- 60	06 04	−26 07	5.1			6.3	1.45	0.37
ESO 5- 4	06 05	−86 37	2			3.3	1.58	0.66
ESO 121- 6	06 07	−61 48	4			5.2	1.58	0.74
ESO 555- 36	06 07	−19 54	5.2			5.0	1.30	1.01

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
UGC 3403	06 10	+71 23	5.2			6.0	1.36	0.52
UGC 3425	06 14	+66 34	4 ^f			3.0	1.40	0.86
UGC 3433	06 15	+00 11	4			4.0	1.31	0.51
UGC 3420	06 16	+75 56	4			3.0	1.41	0.49
IC 2163	06 16	-21 22	5.1			5.0	1.48	0.39
ESO 489- 29	06 17	-27 23	3		SA III	3.7	1.47	0.81
UGCA 127	06 20	-08 27	5.1			6.0	1.59	0.54
NGC 2146A	06 23	+78 31	5.1			5.3	1.48	0.42
UGC 3458	06 25	+64 44	3	3		3.0	1.38	0.60
UGC 3474	06 32	+71 33	4			6.0	1.35	0.98
UGC 3489	06 33	+21 02	4			4.0	1.30	0.84
ESO 58- 13	06 44	-71 27	4			3.0	1.35	0.81
ESO 490- 41	06 46	-26 06	4			-0.4	1.46	0.44
ESO 490- 45	06 46	-26 28	5.1			7.0	1.40	0.45
NGC 2295	06 47	-26 44	3			2.0	1.32	0.54
UGC 3539	06 48	+66 14	4			4.0	1.39	1.04
NGC 2310	06 53	-40 51	2 ^c	2	J B, SA III, S	-2.0	1.64	0.72
UGC 3587	06 53	+19 17	5.1			7 ²	1.45	0.61
ESO 491- 15	07 00	-27 22	3			4.5	1.31	0.69
UGC 3652	07 03	+22 22	5.1			6.0	1.31	0.51
UGC 3697	07 11	+71 50	3			7.0	1.52	1.23
UGC 3717	07 13	+73 50	5.1			4.0	1.32	0.44
ESO 162- 17	07 15	-57 20	5.1			2.5	1.31	0.43
NGC 2369	07 16	-62 20	5.1			1.0	1.55	0.52
NGC 2357	07 17	+23 21	4			6.0	1.55	0.87
ESO 367- 17	07 19	-35 39	5.1			1.5	1.38	0.39
ESO 428- 28	07 23	-30 03	5.2			5.0	1.37	0.70
IC 2202	07 27	-67 34	4			4.3	1.31	0.47
UGC 3855	07 28	+58 30	4			2.0	1.30	0.65
IC 467	07 30	+79 52	5.1			5.0	1.51	0.40
NGC 2410	07 35	+32 49	5.1			3.0	1.39	0.54
ESO 257- 19	07 35	-46 55	5.2			6.2	1.39	0.62
UGC 3909	07 36	+73 42	4			6.0	1.42	0.74
UGC 3937	07 37	+35 36	5.1			5 ²	1.30	0.57
ESO 163- 11	07 38	-55 11	3			3.0	1.42	0.61
UGC 3962	07 40	+13 52	5.1			3.0	1.33	0.45
NGC 2424	07 40	+39 13	1 ^f			3.0	1.58	0.81
UGC 3995	07 44	+29 14	5.1			3 ⁶	1.40	0.35
NGC 2435	07 44	+31 39	4			1.0	1.33	0.63
ESO 123- 23	07 44	-58 09	4			5.7	1.34	0.83
UGC 3984	07 45	+70 01	5.1			3.0	1.30	0.37
ESO 311- 12	07 47	-41 27	2		J B	0.0	1.54	0.91
ESO 560- 13	07 47	-18 44	2+	2+		4.0	1.43	0.79
UGC 4029	07 48	+34 19	4			4.0	1.36	0.68
IC 2207	07 49	+33 57	4 ^f			6.0	1.31	0.83
UGC 4043	07 50	+54 21	4			7.0	1.32	0.95
UGC 4054	07 50	+23 53	4			3.0	1.46	0.67
UGC 4057	07 54	+74 23	4			1.0	1.40	0.52
ESO 35- 18	07 55	-76 24	2			5.3	1.51	0.66
ESO 494- 7	07 56	-24 54	5.2			4.0	1.44	0.66
ESO 209- 9	07 58	-49 51	5.2			6.0	1.79	0.86
UGC 4148	08 00	+42 11	5.5			7.0	1.40	0.87
UGC 4171	08 01	+09 42	4			3.0	1.36	0.86
UGC 4078	08 04	+84 38	4			4.0	1.32	0.69
ESO 494- 22	08 05	-24 48	3			1.0	1.35	0.46
ESO 430- 20	08 07	-28 03	5.6			7.0	1.36	0.48

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log$ D_{25}	(9) $\log$ R_{25}
ESO 89- 12	08 10	−64 56	4			4.0	1.36	0.79
UGC 4257	08 10	+24 53	5.5			6.0	1.33	1.05
ESO 561- 23	08 11	−18 18	4			3.7	1.31	0.81
NGC 2523B	08 12	+73 33	4			3.0	1.33	0.83
UGC 4277	08 13	+52 38	3			6.0	1.59	0.98
IC 2233	08 13	+45 44	3	3		7.0	1.67	1.00
IC 2267	08 18	+24 44	3			6.0	1.32	0.81
NGC 2549	08 18	+57 48	4 ^f		SA II	−2.0	1.59	0.48
MCG -2-22- 5	08 22	−11 25	4			4.0	1.31	0.67
NGC 2590	08 25	−00 35	4			3.5	1.35	0.49
UGC 4416	08 27	+22 52	5.1			3.0	1.35	0.37
ESO 562- 7	08 28	−21 44	5.1			2.0	1.38	0.48
NGC 2613	08 33	−22 58	3			3.0	1.86	0.61
NGC 2612	08 33	−13 10	4			−3.0	1.43	0.68
UGC 4491	08 35	+01 43	5.1			1.0	1.38	0.45
ESO 562- 23	08 36	−20 28	3			−0.5	1.34	0.64
NGC 2591	08 37	+78 01	4			5.5	1.48	0.69
NGC 2620	08 37	+24 56	4			4 ¹	1.31	0.59
UGC 4514	08 39	+53 27	5.1			6.0	1.33	0.36
ESO 432- 2	08 40	−32 02	5.2			5.0	1.33	1.01
NGC 2644	08 41	+04 58	5.1			4 ²	1.33	0.41
NGC 2648	08 42	+14 17	4			1.0	1.51	0.47
ESO 563- 14	08 42	−20 03	4			6.7	1.36	0.68
UGC 4550	08 43	+13 05	3			3.0	1.43	0.81
UGC 4551	08 44	+49 47	4			−2.0	1.31	0.48
UGC 4559	08 44	+30 07	4 ^f			2.0	1.51	0.82
ESO 563- 21	08 47	−20 02	4			4.0	1.48	0.84
NGC 2654	08 49	+60 13	1	1	J P, SA I, S	2.0	1.63	0.73
ESO 371- 20	08 50	−34 32	5.1			5.0	1.50	0.47
MCG -3-23- 10	08 51	−17 33	5.1			6.9	1.55	0.53
UGC 4640	08 51	−02 08	5.1			5.3	1.51	0.46
NGC 2683	08 52	+33 25	2	2	SA II, S	3.0	1.97	0.63
UGC 4623	08 53	+76 29	4			6.0	1.55	0.61
ESO 371- 26	08 54	−32 56	3			−1.7	1.44	0.69
NGC 2713	08 57	+02 55	5.1			2.0	1.56	0.38
ESO 60- 19	08 57	−69 03	5.1			7.0	1.50	0.41
MCG -1-23- 13	08 58	−06 11	3			4.0	1.30	0.41
ESO 90- 12	08 58	−66 43	4			−2.0	1.31	0.60
UGC 4719	09 00	+50 40	4			5.0	1.36	0.92
NGC 2788A*	09 02	−68 13	1 ^b	1	J P, SDB	2.5	1.45	0.84
ESO 36- 19	09 06	−75 48	5.1			3.9	1.30	0.42
NGC 2715	09 08	+78 05	5.1			5.0	1.69	0.47
NGC 2770	09 09	+33 07	4			5.0	1.58	0.52
MCG -1-24- 1	09 10	−08 54	3			2.8	1.63	0.62
NGC 2781	09 11	−14 49	5.1			−1.0	1.48	0.30
UGC 4824	09 11	+51 15	4			7.0	1.39	0.59
ESO 564- 27	09 11	−20 07	4 ^c			6.3	1.61	1.09
ESO 433- 8	09 12	−30 54	4			−2.0	1.30	0.79
NGC 2784	09 12	−24 10	4			−2.0	1.74	0.39
NGC 2732	09 13	+79 11	4		SA III	−2.0	1.32	0.42
NGC 2748	09 13	+76 28	3			4.0	1.48	0.42
UGC 4869	09 14	+30 08	4			−2.0	1.38	0.55
ESO 564- 32	09 14	−21 58	3			4.0	1.31	0.75
NGC 2811	09 16	−16 18	5.1			1.0	1.40	0.46
NGC 2815	09 16	−23 38	5.1			3.0	1.54	0.49
NGC 2821	09 16	−26 49	3			4.0	1.31	0.66

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log$ D_{25}	(9) $\log$ R_{25}
NGC 2798	09 17	+42 00	5.1			1.0	1.41	0.42
UGC 4906	09 17	+52 59	4		SA III	1.0	1.30	0.59
ESO 433- 12	09 18	-32 28	5.1			3.7	1.36	0.69
IC 2461	09 19	+37 11	4			3.0	1.37	0.75
ESO 372- 16	09 21	-33 11	3			3.2	1.38	0.74
NGC 2820	09 21	+64 15	3 ^f			5.0	1.61	0.92
NGC 2841	09 22	+50 58	5.1			3.0	1.91	0.36
IC 2469	09 23	-32 26	1		S	2.0	1.67	0.66
NGC 2862	09 24	+26 46	1	1		3 ⁶	1.40	0.66
ESO 372- 23	09 25	-37 10	2			4.5	1.30	0.95
NGC 2874	09 25	+11 25	5.1			4.0	1.38	0.51
UGC 5020	09 26	+34 39	4			6.0	1.33	0.68
NGC 2870	09 27	+57 22	4			4.0	1.39	0.58
ESO 373- 8	09 33	-33 01	5.2			6.0	1.76	0.82
NGC 2921	09 34	-20 55	5.1			1.0	1.45	0.44
MCG -1-25- 5	09 35	-07 43	3			-1.0	1.32	0.49
NGC 2939	09 38	+09 31	3	3		4.0	1.39	0.45
UGC 5173	09 41	+11 24	4			3.0	1.36	0.92
NGC 2966	09 42	+04 40	5.1			5 ²	1.35	0.41
MCG -1-25- 26	09 42	-06 44	2+			3.0	1.34	0.75
NGC 2992	09 45	-14 19	2			1.0	1.55	0.51
UGC 5228	09 46	+01 40	5.1			5.0	1.39	0.53
UGC 5210	09 46	+68 58	5.5			6.0	1.35	1.06
ESO 37- 5	09 46	-74 35	5.1			3.0	1.47	0.57
UGC 5203	09 47	+79 48	5.5			7.0	1.36	1.10
ESO 499- 5	09 47	-24 50	5.2			4.5	1.40	0.56
UGC 5249	09 47	+02 37	5.1			7.0	1.38	0.51
NGC 3003	09 48	+33 25	5.4			4.0	1.76	0.63
IC 2511	09 49	-32 50	2			1.4	1.46	0.70
NGC 3024	09 50	+12 45	4			5.0	1.32	0.66
ESO 374- 3	09 51	-33 04	5.1			5.5	1.34	0.46
NGC 3044	09 53	+01 34	2			5.0	1.69	0.84
NGC 3060	09 56	+16 49	4			3.0	1.35	0.60
UGC 5341	09 56	+20 38	4			6.0	1.47	1.13
IC 2526	09 57	-32 15	4			-2.0	1.32	0.48
ESO 435- 14	09 57	-28 30	2			5.0	1.41	0.92
NGC 3067	09 58	+32 22	5.1			2.0	1.39	0.42
ESO 435- 19	09 59	-30 14	3			5.0	1.50	0.96
IC 2531	09 59	-29 36	1 ^c		J P, SA I, S	5.3	1.84	1.09
NGC 3079	10 01	+55 40	1		S	5.0 ³	1.90	0.74
NGC 3098	10 02	+24 42	3 ^f	3	SA I	-2.0	1.36	0.57
MCG -1-26- 12	10 02	-06 00	4			7.0	1.45	1.02
ESO 316- 18	10 02	-42 05	4			4.7	1.38	0.87
ESO 499- 37	10 03	-27 01	5.1			7.0	1.51	0.38
ESO 316- 21	10 04	-41 24	4			3.0	1.41	0.96
NGC 3113	10 04	-28 26	5.1			7.3	1.52	0.44
NGC 3115	10 05	-07 43	2		SA II	-3.0	1.86	0.47
NGC 3118	10 07	+33 01	4			4.0	1.40	0.83
UGC 5459	10 08	+53 04	3			5.3	1.68	0.81
NGC 3126	10 08	+31 51	4			3.0	1.45	0.71
NGC 3131	10 08	+18 13	5.1			3.0	1.38	0.54
IC 2554	10 08	-67 01	5.3			4.0	1.49	0.38
NGC 3137	10 09	-29 03	5.1			6.3	1.80	0.45
NGC 3157	10 11	-31 38	3			4.5	1.39	0.67
UGC 5495	10 11	+16 26	4			6.0	1.48	0.88
UGC 5499	10 12	+27 51	3			3.0	1.44	0.68

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
ESO 263- 15	10 12	-47 17	4			6.0	1.46	0.94
ESO 436- 1	10 12	-27 50	5.2			4.3	1.46	0.92
NGC 3153	10 12	+12 40	5.1			6.0	1.33	0.36
ESO 263- 18	10 13	-43 43	4			4.0	1.31	0.77
ESO 567- 26	10 14	-21 58	4			4.0	1.30	0.72
NGC 3175	10 14	-28 52	5.1			1.0	1.70	0.57
ESO 263- 23	10 14	-43 37	4			0.0	1.34	0.58
UGC 5537	10 15	+07 19	5.5			7.0	1.40	1.15
NGC 3187	10 17	+21 52	5.1			5.0	1.47	0.37
NGC 3190	10 18	+21 49	4			1.0	1.64	0.45
NGC 3200	10 18	-17 58	5.1			4.5	1.62	0.51
NGC 3203	10 19	-26 41	3	3	SA III	-1.0	1.46	0.69
NGC 3198	10 19	+45 33	5.1			5.0	1.93	0.41
ESO 263- 29	10 21	-46 00	2			-0.6	1.34	0.61
NGC 3221	10 22	+21 34	4			6.0	1.51	0.68
NGC 3230	10 23	+12 33	5.1			-2.0	1.36	0.30
UGC 5646	10 25	+14 21	4			5 ¹	1.44	0.56
NGC 3245A	10 27	+28 38	3			3.0	1.52	0.99
NGC 3250B	10 27	-40 26	1 ^b			1.0	1.36	0.59
NGC 3248	10 27	+22 50	4			-2.0	1.40	0.34
ESO 436- 27	10 28	-31 36	5.1			-2.0	1.37	0.30
UGC 5687	10 29	+06 07	4			6.0	1.31	0.96
NGC 3263	10 29	-44 06	5.2			6.0	1.71	0.56
NGC 3251	10 29	+26 05	4			4 ¹	1.30	0.70
NGC 3254	10 29	+29 29	4			4.0	1.70	0.50
IC 2584	10 29	-34 54	4			-2.0	1.30	0.65
NGC 3269	10 29	-35 13	5.1			-1.0	1.40	0.34
UGC 5708	10 31	+04 28	5.4			7.0	1.54	0.75
NGC 3270	10 31	+24 52	4 ^f			3.0	1.50	0.58
ESO 436- 34	10 32	-28 36	4 ^d			3.0	1.32	0.74
ESO 317- 46	10 33	-38 58	4			7.0	1.36	0.85
NGC 3289	10 34	-35 19	5.1			-1.0	1.35	0.60
NGC 3281D	10 34	-34 24	3			6.7	1.31	0.70
NGC 3252	10 34	+73 45	5.1			6.7	1.30	0.50
NGC 3279	10 34	+11 11	4			7.0	1.46	0.92
UGC 5751	10 35	+21 02	5.1			4 ²	1.31	0.65
IC 624	10 36	-08 20	4			1.0	1.43	0.55
ESO 437- 14	10 36	-32 20	4			2.4	1.41	0.53
NGC 3301	10 36	+21 52	4			0.0	1.55	0.54
ESO 437- 15	10 36	-28 10	4			-1.7	1.33	0.65
NGC 3312	10 37	-27 33	5.1			3.0	1.52	0.42
UGC 5757	10 38	+79 22	5.1			1.0	1.32	0.40
ESO 437- 30	10 39	-30 17	3			3.9	1.49	0.72
NGC 3320	10 39	+47 23	5.1			6.0	1.34	0.35
NGC 3333	10 39	-36 02	4			3.8	1.30	0.75
NGC 3347A	10 40	-36 24	5.1			5.8	1.30	0.43
ESO 501- 80	10 42	-23 56	4			5.0	1.37	0.64
ESO 318- 4	10 43	-38 15	2			5.0	1.43	0.76
NGC 3365	10 46	+01 48	4			6.0	1.65	0.76
ESO 318- 13	10 47	-38 51	5.2			7.0	1.35	0.85
UGC 5897	10 47	+11 04	5.1			5.0	1.42	0.44
NGC 3390	10 48	-31 31	2 ^c		J B, S	3.0	1.55	0.79
NGC 3384	10 48	+12 37	5.1			-3.0	1.74	0.34
UGC 5904	10 48	+66 21	2	3		3.0	1.30	0.79
ESO 264- 43	10 48	-45 25	3			3.0	1.30	0.64
NGC 3404	10 50	-12 06	3			1.8	1.33	0.64

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log$ D_{25}	(9) $\log$ R_{25}
NGC 3413	10 51	+32 46	4			-2.0	1.34	0.38
ESO 569- 14	10 51	-19 53	3			6.2	1.52	0.85
UGC 5974	10 51	+04 34	5.1			6.0	1.31	0.40
PGC 32573	10 51	-17 07	5.1			1.7	1.34	0.39
NGC 3424	10 51	+32 54	5.1			3.0	1.45	0.57
NGC 3437	10 52	+22 56	5.1			5.0	1.40	0.49
ESO 264- 48	10 52	-45 40	5.2			5.0	1.31	0.51
NGC 3449	10 52	-32 55	4			2.0	1.52	0.54
NGC 3440	10 53	+57 07	4			3.0	1.32	0.62
NGC 3403	10 53	+73 41	5.1			4.0	1.48	0.41
ESO 376- 26	10 54	-33 07	5.1			-1.0	1.32	0.31
NGC 3454	10 54	+17 20	3			5.0	1.32	0.71
MCG -2-28- 26	10 58	-15 31	5.1			?	1.30	0.91
NGC 3478	10 59	+46 07	5.1			3.5	1.42	0.35
UGC 6080	11 00	+61 19	4			7.0	1.33	1.11
ESO 265- 3	11 01	-44 02	5.1			6.3	1.43	0.36
NGC 3495	11 01	+03 37	5.4			7.0	1.69	0.61
UGC 6112	11 02	+16 44	5.1			7.0	1.40	0.49
NGC 3501	11 02	+17 59	3			6.0	1.59	0.88
NGC 3511	11 03	-23 05	5.1			5.0	1.76	0.45
NGC 3509	11 04	+04 49	5.1			4.0	1.33	0.35
ESO 377- 10	11 06	-37 39	3			2.7	1.41	0.59
NGC 3533	11 07	-37 10	5.2			1.8	1.45	0.64
ESO 265- 7	11 07	-46 31	5.1			5.5	1.56	0.48
NGC 3568	11 10	-37 26	5.1			5.0	1.40	0.50
ESO 377- 21	11 10	-35 58	5.1			3.1	1.31	0.42
NGC 3549	11 10	+53 23	5.1			5.0	1.51	0.44
NGC 3573	11 11	-36 52	3			0.0	1.56	0.57
NGC 3571	11 11	-18 17	5.1			0.7	1.48	0.46
NGC 3556	11 11	+55 40	5.2			6.0	1.94	0.59
NGC 3593	11 14	+12 49	4		SA III	0.0	1.72	0.43
ESO 438- 15	11 15	-28 23	5.1			4.1	1.30	0.35
NGC 3600	11 15	+41 35	4			1.0	1.61	0.68
ESO 377- 31	11 16	-33 57	5.1			3.8	1.31	0.37
NGC 3620	11 16	-76 12	5.1			1.7	1.44	0.38
NGC 3607	11 16	+18 03	5.1			-2.0	1.69	0.30
ESO 377- 34	11 17	-34 57	4			4.0	1.34	0.55
NGC 3623	11 18	+13 05	5.1			1.0	1.99	0.53
NGC 3638	11 20	-08 06	4			3.0	1.34	0.49
NGC 3628	11 20	+13 35	1 ^e	1	SA II, SDB	3.0	2.17	0.70
NGC 3625	11 20	+57 46	4			3.0	1.30	0.49
UGC 6355	11 20	+31 13	5.5			7.0	1.30	0.77
UGC 6378	11 22	+69 38	2			7.0	1.44	1.04
NGC 3652	11 22	+37 46	5.3			6.0	1.30	0.48
UGC 6390	11 22	+64 04	3			7.0	1.32	0.84
UGC 6415	11 24	+27 01	4			2.0	1.32	0.69
NGC 3666	11 24	+11 20	4			5.0	1.64	0.56
NGC 3669	11 25	+57 43	2	2		6.0	1.35	0.61
ESO 170- 4	11 26	-54 15	5.6			5.0	1.36	0.73
NGC 3693	11 28	-13 11	5.1			2.7	1.51	0.68
NGC 3692	11 28	+09 24	4			3.0	1.50	0.65
NGC 3697	11 28	+20 47	5.1			3.0	1.37	0.50
UGC 6483	11 29	+17 13	3			6.0	1.35	1.06
UGC 6497	11 30	+38 37	5.5			7.0	1.36	0.96
NGC 3705	11 30	+09 16	5.1			2.0	1.69	0.38
UGC 6509	11 31	+23 06	5.5			7.0	1.32	1.12

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log$ D_{25}	(9) $\log$ R_{25}
NGC 3717	11 31	−30 18	2+			3.0	1.78	0.73
UGC 6534	11 33	+63 16	5.1			7.0	1.40	0.61
NGC 3749	11 35	−37 59	3			1.0	1.50	0.60
NGC 3735	11 35	+70 32	3			5.0	1.62	0.70
NGC 3755	11 36	+36 24	5.1			5.0	1.50	0.36
UGC 6594	11 37	+16 33	4			7.0	1.41	0.84
NGC 3769	11 37	+47 53	5.1			3.0	1.49	0.50
MCG -3-30- 3	11 37	−17 14	2			5.0	1.32	0.83
UGC 6603	11 38	+35 12	4			6.0	1.34	0.67
UGC 6610	11 38	+33 48	4			6.0	1.32	0.73
NGC 3788	11 39	+31 55	3			2.0	1.33	0.51
ESO 216- 31	11 40	−48 57	3			2.0	1.31	0.64
NGC 3795	11 40	+58 36	4			4 ¹	1.33	0.59
NGC 3800	11 40	+15 20	5.1			3.0	1.31	0.54
UGC 6667	11 42	+51 36	3			6.0	1.53	0.89
NGC 3831	11 43	−12 52	2			−0.5	1.43	0.68
UGC 6686	11 43	+16 29	4			3.0	1.42	0.92
IC 724	11 43	+08 56	4			1.0	1.37	0.40
NGC 3863	11 45	+08 28	4			4.0	1.45	0.68
MCG -2-30- 27	11 45	−10 06	5.1			6.5	1.44	0.55
NGC 3877	11 46	+47 29	5.1			5.0	1.74	0.63
NGC 3885	11 46	−27 55	4			0.0	1.38	0.40
UGC 6774	11 48	+55 02	4			6.0	1.31	0.91
UGC 6780	11 48	−02 01	5.1			6.7	1.51	0.52
UGC 6792	11 49	+39 46	3			6.0	1.44	0.83
ESO 320- 26	11 49	−38 47	5.1			3.0	1.38	0.43
UGC 6802	11 50	+51 52	4			6.0	1.33	1.12
UGC 6806	11 50	+25 57	5.1			1 ²	1.30	0.48
NGC 3917	11 50	+51 49	4			6.0	1.71	0.61
NGC 3936	11 52	−26 54	2			4.0	1.59	0.79
ESO 379- 6	11 53	−36 38	4			4.7	1.40	0.81
ESO 440- 27	11 53	−28 33	3			7.0	1.64	0.84
IC 2974	11 53	−05 10	4			5.0	1.34	0.68
NGC 3955	11 53	−23 09	5.1			0.0	1.46	0.49
NGC 3956	11 54	−20 34	5.1			5.0	1.53	0.53
NGC 3957	11 54	−19 34	3 ^c		SA I, S	−1.0	1.49	0.66
ESO 320- 31	11 54	−39 51	2+			4.7	1.42	0.94
NGC 3972	11 55	+55 19	4			4.0	1.59	0.56
NGC 3976	11 55	+06 44	5.1			3.0	1.58	0.50
NGC 3981	11 56	−19 53	5.1			4.0	1.72	0.35
NGC 3986	11 56	+32 01	1	1		−2.0	1.49	0.65
NGC 3987	11 57	+25 11	1	1		3.0	1.35	0.73
NGC 4013	11 58	+43 56	1		J B, SA I, S	3.0	1.72	0.71
NGC 4010	11 58	+47 15	5.4			6.7	1.63	0.73
UGC 6968	11 58	+28 17	5.1			5 ⁶	1.45	0.44
IC 750	11 58	+42 43	5.1			2.0	1.42	0.35
NGC 4020	11 58	+30 24	5.1			7.3	1.32	0.35
NGC 4026	11 59	+50 57	4 ^a	4	SA I	−2.0	1.72	0.61
IC 755	12 01	+14 06	2	3		3.0	1.38	0.88
NGC 4036	12 01	+61 53	4			−3.0	1.63	0.40
NGC 4062	12 04	+31 53	5.1			5.0	1.61	0.37
NGC 4064	12 04	+18 26	5.1			1.0	1.64	0.40
NGC 4085	12 05	+50 21	3			5.0	1.45	0.55
NGC 4088	12 05	+50 32	5.1			4.0	1.76	0.41
MCG 5-29- 11	12 05	+31 03	5.3			−3.0	1.42	0.41
IC 2995	12 05	−27 56	5.1			5.0	1.51	0.52

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
NGC 4094	12 05	−14 31	5.1			5.5	1.62	0.45
UGC 7086	12 05	+77 30	4			3.0	1.43	0.87
NGC 4096	12 06	+47 28	5.4			5.0	1.82	0.57
NGC 4100	12 06	+49 34	5.1			4.0	1.73	0.48
NGC 4111	12 07	+43 04	3 ^a		J P, SA I	−1.0	1.66	0.67
IC 3005	12 07	−30 01	4			5.5	1.37	0.78
NGC 4124	12 08	+10 22	4			−1.0	1.63	0.48
ESO 441- 7	12 08	−32 00	4			4.0	1.32	0.84
NGC 4128	12 08	+68 46	4			−2.0	1.42	0.48
NGC 4129	12 08	−09 02	3		SA II	2.0	1.37	0.59
IC 3015	12 09	−31 31	3			3.5	1.46	0.63
NGC 4134	12 09	+29 10	5.1			3.0	1.34	0.40
UGC 7153	12 09	+62 16	4			6.0	1.30	0.83
NGC 4144	12 09	+46 27	5.2			6.0	1.78	0.66
IC 764	12 10	−29 44	5.1			5.0	1.68	0.47
UGC 7170	12 10	+18 49	3 ^d			6.0	1.48	1.13
NGC 4157	12 11	+50 29	3			3.0	1.83	0.70
NGC 4173	12 12	+29 12	5.4			7.0	1.70	0.84
NGC 4179	12 12	+01 18	4			−2.0	1.60	0.55
NGC 4183	12 13	+43 41	4			6.0	1.72	0.83
NGC 4192	12 13	+14 53	2			2.0	1.99	0.55
NGC 4197	12 14	+05 48	4			7.0	1.53	0.78
IC 3061	12 15	+14 01	4			5.3	1.35	0.71
ESO 441- 22	12 15	−30 45	4			3.7	1.32	0.57
NGC 4206	12 15	+13 01	4			4.0	1.79	0.72
ESO 321- 16	12 15	−38 08	5.1			5.5	1.41	0.54
UGC 7271	12 15	+43 26	5.1			7.0	1.31	0.53
ESO 380- 6	12 15	−35 37	5.1			2.7	1.56	0.37
NGC 4217	12 15	+47 05	4	4		3.0	1.72	0.53
NGC 4216	12 15	+13 08	2			3.0	1.91	0.66
NGC 4220	12 16	+47 53	4			−1.0	1.59	0.45
NGC 4222	12 16	+13 18	4			7.0	1.52	0.86
NGC 4219	12 16	−43 19	5.1			4.0	1.63	0.53
NGC 4224	12 16	+07 27	3			1.0	1.41	0.40
NGC 4233	12 17	+07 37	5.1			−1.5	1.38	0.34
NGC 4235	12 17	+07 11	3		S	1.0	1.62	0.65
NGC 4244	12 17	+37 48	3			6.0	2.22	0.94
UGC 7321	12 17	+22 32	4			7.0	1.74	1.19
NGC 4251	12 18	+28 10	4			−2.0	1.56	0.39
NGC 4256	12 18	+65 53	4			3.0	1.65	0.77
NGC 4258	12 18	+47 18	5.1			4.0	2.27	0.41
NGC 4266	12 19	+05 32	4			1.0	1.30	0.68
NGC 4270	12 19	+05 27	5.1			−2.0	1.31	0.35
NGC 4274	12 19	+29 36	5.1			2.0	1.83	0.43
ESO 506- 2	12 20	−26 04	4			4.3	1.38	0.96
NGC 4281	12 20	+05 23	5.1			−1.0	1.48	0.31
NGC 4289	12 21	+03 43	3			5.5	1.59	1.01
NGC 4294	12 21	+11 30	5.1			6.0	1.51	0.42
NGC 4302	12 21	+14 36	2+	2		5.0	1.74	0.74
ESO 380- 19	12 22	−35 47	4			6.0	1.49	0.82
NGC 4307	12 22	+09 02	4			3.0	1.56	0.67
NGC 4312	12 22	+15 32	3			1.5	1.66	0.61
NGC 4313	12 22	+11 48	3			2.0	1.60	0.59
NGC 4316	12 22	+09 19	4			6.0	1.39	0.70
NGC 4324	12 23	+05 14	5.1			−1.0	1.44	0.35
NGC 4330	12 23	+11 22	3			6.0	1.65	0.70

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log$ D_{25}	(9) $\log$ R_{25}
NGC 4338	12 23	+28 53	4			7.0	1.34	1.01
NGC 4346	12 23	+46 59	4		SA III	-2.0	1.52	0.42
NGC 4343	12 23	+06 57	4			3.0	1.40	0.54
IC 3253	12 23	-34 37	5.1			4.8	1.45	0.39
NGC 4348	12 23	-03 26	3			4.0	1.51	0.64
NGC 4350	12 23	+16 41	4		SA III	-2.0	1.48	0.32
NGC 4357	12 23	+48 46	5.1			4.0	1.56	0.45
NGC 4352	12 24	+11 13	5.1			-2.0	1.33	0.35
NGC 4359	12 24	+31 31	3			5.0	1.55	0.65
NGC 4356	12 24	+08 32	3			6.0	1.45	0.71
NGC 4373A	12 25	-39 19	4			-1.5	1.41	0.49
IC 3322A	12 25	+07 13	2			6.0	1.54	0.89
NGC 4388	12 25	+12 39	2+		SA II, S	3.0	1.75	0.64
IC 3322	12 25	+07 33	4			6.0	1.37	0.69
UGC 7522	12 25	+03 25	3			6.0	1.45	0.95
NGC 4396	12 25	+15 40	5.1			7.3	1.52	0.53
NGC 4402	12 26	+13 06	2			3.0	1.59	0.55
NGC 4417	12 26	+09 35	4			-2.0	1.53	0.41
NGC 4419	12 26	+15 02	4			1.0	1.52	0.47
NGC 4425	12 27	+12 44	3		SA II	-1.0	1.47	0.47
NGC 4429	12 27	+11 06	2			-1.0	1.75	0.34
NGC 4433	12 27	-08 16	5.1			2.0	1.34	0.36
NGC 4438	12 27	+13 00	5.1			0.0	1.93	0.43
NGC 4442	12 28	+09 48	3			-2.0	1.66	0.41
NGC 4445	12 28	+09 26	4			2.0	1.42	0.73
NGC 4448	12 28	+28 37	5.1			2.0	1.59	0.44
NGC 4452	12 28	+11 45	3		SA II	-2.0	1.44	0.66
IC 3392	12 28	+15 00	5.1			2.5	1.36	0.37
NGC 4455	12 28	+22 49	5.2			7.3	1.44	0.54
NGC 4460	12 28	+44 51	4 ^a		SA III	-1.0	1.60	0.53
UGC 7607	12 28	+04 17	5.5			7.0	1.34	1.25
NGC 4461	12 29	+13 11	4			-1.0	1.55	0.39
UGC 7617	12 29	+44 39	4			7.0	1.39	1.00
NGC 4462	12 29	-23 09	5.1			2.0	1.51	0.42
NGC 4469	12 29	+08 45	1 ^d	1	J P, SA I, S	0.0	1.58	0.47
UGC 7644	12 30	+03 44	3			7.0	1.34	0.63
NGC 4488	12 30	+08 21	3			0.0	1.57	0.39
NGC 4497	12 31	+11 37	5.1			-1.0	1.31	0.34
ESO 218- 8	12 31	-51 44	4			5 ⁷	1.41	1.22
NGC 4503	12 32	+11 10	5.1			-3.0	1.55	0.33
IC 3474	12 32	+02 39	5.5			7.0	1.36	0.90
NGC 4517	12 32	+00 06	4			6.0	2.02	0.83
UGC 7699	12 32	+37 37	5.1			6.0	1.58	0.57
NGC 4521	12 32	+63 56	4			0.0	1.40	0.66
UGC 7697	12 32	+20 10	4			6.0	1.32	0.76
NGC 4522	12 33	+09 10	5.1			6.0	1.57	0.57
NGC 4526	12 34	+07 42	4			-2.0	1.86	0.48
NGC 4527	12 34	+02 39	3			4.0	1.79	0.47
NGC 4533	12 34	+02 19	4			7.1	1.32	0.69
NGC 4536	12 34	+02 11	5.1			4.0	1.88	0.37
NGC 4539	12 34	+18 12	5.1			0.5	1.52	0.39
NGC 4546	12 35	-03 47	5.1			-3.0	1.52	0.36
NGC 4550	12 35	+12 13	4			-1.5	1.52	0.55
NGC 4544	12 35	+03 02	4			0.0	1.30	0.50
NGC 4559	12 35	+27 57	5.1			6.0	2.03	0.39
NGC 4553	12 36	-39 26	5.1			-1.0	1.33	0.32

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
NGC 4565	12 36	+25 59	2 ^a		J P, SA II, S	3.0	2.20	0.87
UGC 7774	12 36	+40 00	5.4			7.0	1.56	0.92
NGC 4568	12 36	+11 14	5.1			4.0	1.66	0.36
NGC 4570	12 36	+07 14	4			-2.0	1.58	0.52
ESO 442- 13	12 37	-28 29	5.1			5.7	1.47	0.46
NGC 4586	12 38	+04 19	2		SDB	1.0	1.60	0.47
IC 3608	12 38	+10 28	4			3.0	1.49	0.88
ESO 322- 42	12 38	-42 12	5.2			5.5	1.31	0.55
NGC 4594	12 39	-11 37	4		SA I	1.0	1.94	0.39
NGC 4605	12 40	+61 36	5.1			5.0	1.76	0.42
NGC 4602	12 40	-05 07	5.1			4.0	1.53	0.46
UGC 7848	12 40	+63 30	5.1			6.0	1.35	0.39
NGC 4617	12 41	+50 23	3			3.0	1.47	0.74
NGC 4607	12 41	+11 53	4			3.0	1.46	0.64
NGC 4631	12 42	+32 32	3			7.0	2.19	0.76
NGC 4623	12 42	+07 40	4			-0.5	1.35	0.49
NGC 4632	12 42	-00 04	5.1			5.0	1.49	0.42
NGC 4634	12 42	+14 17	4	4		6.0	1.41	0.58
UGC 7883	12 42	-01 13	5.4			6.2	1.42	0.58
NGC 4645A	12 43	-41 21	4			-2.0	1.47	0.56
NGC 4658	12 44	-10 05	5.1			4.0	1.32	0.35
IC 3718	12 44	+12 21	5.1			6 ²	1.43	0.45
NGC 4666	12 45	-00 27	2			5.0	1.66	0.55
ESO 507- 7	12 45	-26 14	4			4.0	1.38	0.90
UGC 7941	12 46	+64 33	4			7.0	1.47	0.70
NGC 4676A	12 46	+30 44	5.1			-2.0	1.36	0.51
NGC 4676B	12 46	+30 43	5.1			0.0	1.34	0.44
NGC 4672	12 46	-41 42	4			1.4	1.31	0.56
NGC 4686	12 46	+54 32	4			1.0	1.31	0.55
NGC 4693	12 47	+71 10	4			7.0	1.39	0.71
NGC 4684	12 47	-02 43	5.1			-1.0	1.46	0.44
NGC 4679	12 47	-39 34	5.1			4.7	1.38	0.40
NGC 4694	12 48	+10 59	5.1			-2.0	1.50	0.32
IC 3799	12 48	-14 23	3			7.0	1.37	0.93
NGC 4700	12 49	-11 24	3			5.0	1.48	0.73
NGC 4703	12 49	-09 06	4			3.0	1.47	0.71
NGC 4705	12 49	-05 11	5.1			4.0	1.48	0.46
NGC 4712	12 49	+25 28	5.1			4.0	1.40	0.35
NGC 4710	12 49	+15 09	2+		SA II, S	-1.0	1.69	0.62
UGC 7982	12 49	+02 51	4			4 ¹	1.53	0.66
UGC 7993	12 50	+52 07	4			6.0	1.33	0.90
NGC 4738	12 51	+28 47	4			6.0	1.32	0.83
NGC 4747	12 51	+25 46	5.1			6.0	1.54	0.47
NGC 4746	12 51	+12 05	5.1			3.0	1.58	0.52
ESO 442- 26	12 52	-29 50	4			-2.3 ⁵	1.43	0.53
MCG -1-33- 32	12 52	-09 45	4			6.4	1.37	0.95
NGC 4751	12 52	-42 39	3			-3.0	1.48	0.42
NGC 4762	12 52	+11 13	3	4	SA I	-2.0	1.94	0.72
NGC 4771	12 53	+01 16	4			7.0	1.59	0.67
NGC 4781	12 54	-10 32	5.1			7.0	1.54	0.35
MCG -2-33- 52	12 54	-11 36	4			5 ⁶	1.30	0.91
UGC 8032	12 54	+13 13	2			-2 ⁴	1.44	0.59
IC 3881	12 54	+19 10	4			6.0	1.57	0.62
ESO 323- 42	12 55	-40 58	5.1			4 ⁶	1.35	0.47
NGC 4808	12 55	+04 18	5.1			6.0	1.44	0.38
NGC 4818	12 56	-08 31	5.1			1.7	1.63	0.45

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
UGC 8067	12 57	-01 42	4			3.0	1.32	0.73
NGC 4835A	12 57	-46 22	2 ^d		SDB	6.0	1.41	0.80
MCG -1-33- 60	12 57	-09 38	4			6.7	1.49	0.98
NGC 4843	12 58	-03 37	2			0.0	1.32	0.64
NGC 4845	12 58	+01 34	1		SA II, S	2.0	1.70	0.58
NGC 4835	12 58	-46 15	5.2			4.0	1.60	0.67
UGC 8085	12 58	+14 33	4			6.0	1.40	0.63
NGC 4856	12 59	-15 2	5.1			0.0	1.63	0.56
NGC 4866	12 59	+14 10	3			-1.0	1.80	0.67
MCG -1-33- 71	13 01	-08 20	1 ^d			5.0	1.42	0.75
UGC 8146	13 02	+58 41	3			7.0	1.54	0.98
ESO 219- 21	13 02	-50 20	5.1			7.0	1.70	0.64
MCG -3-33- 28	13 02	-17 40	2			5.0	1.41	1.08
ESO 443- 42	13 03	-29 49	1 ^b	1	J P, SDB	3.0	1.44	0.8 2
PGC 45134	13 03	-16 33	4			5.0	1.30	0.72
NGC 4951	13 05	-06 29	5.1			6.0	1.52	0.45
NGC 4945	13 05	-49 28	2			6.0	2.30	0.72
NGC 4958	13 05	-08 01	4		SA II, S	-2.0	1.61	0.51
ESO 508- 11	13 07	-22 51	5.2			7.0	1.48	0.79
UGC 8246	13 10	+34 10	3			6.0	1.54	0.73
ESO 576- 3	13 10	-21 44	5.1			6.7	1.30	0.46
ESO 269- 61	13 11	-44 35	4			3.0	1.40	0.56
IC 4213	13 12	+35 40	3			6.0	1.40	0.73
NGC 5023	13 12	+44 02	3			5.5	1.78	0.89
MCG -3-34- 14	13 12	-17 32	5.1			4.7	1.40	0.51
NGC 5025	13 12	+31 48	4			3.0	1.31	0.52
ESO 443- 83	13 12	-32 41	5.2			6.9	1.53	0.60
ESO 576- 11	13 13	-19 58	2			6.4	1.47	0.80
UGC 8304	13 13	+50 36	4			3.0	1.49	0.66
NGC 5022	13 13	-19 32	3			3.0	1.38	0.83
ESO 219- 41	13 13	-49 28	5.2			2.2	1.45	0.51
NGC 5037	13 14	-16 35	4			1.0	1.35	0.48
NGC 5047	13 15	-16 31	1 ^b			-2.0	1.45	0.69
MCG -3-34- 41	13 17	-16 15	2			5.0	1.36	0.92
ESO 382- 31	13 17	-37 16	2			6.5	1.37	0.80
NGC 5062	13 18	-35 27	5.1			-2.0	1.35	0.49
NGC 5081	13 19	+28 30	5.1			3.0	1.35	0.44
NGC 5073	13 19	-14 50	1			5.0	1.53	0.69
ESO 382- 41	13 19	-35 06	5.2			5.0	1.31	1.07
NGC 5084	13 20	-21 49	4			-2.0	1.97	0.73
NGC 5088	13 20	-12 34	5.1			3.7	1.42	0.52
ESO 576- 40	13 20	-22 03	4			7.0	1.32	0.72
ESO 40- 7	13 21	-77 32	5.2			5.3	1.42	0.81
NGC 5102	13 21	-36 37	4			-3.0	1.94	0.49
NGC 5116	13 22	+26 58	5.1			5.0	1.31	0.44
NGC 5131	13 23	+30 59	4			1.0	1.33	0.79
ESO 382- 58	13 25	-33 39	3			3.7	1.37	0.78
ESO 324- 23	13 27	-38 10	5.2			6.5	1.58	0.73
NGC 5152	13 27	-29 37	5.1			3.0	1.30	0.49
ESO 509- 19	13 27	-25 51	4			4.0	1.38	1.12
NGC 5169	13 28	+46 40	5.1			3.0	1.37	0.40
NGC 5166	13 28	+32 01	2+	1		3.0	1.36	0.80
IC 4263	13 28	+46 55	5.1			6.7	1.31	0.68
ESO 324- 29	13 29	-41 59	4			-0.6	1.38	0.48
NGC 5161	13 29	-33 10	5.1			5.0	1.75	0.41
ESO 383- 5	13 29	-34 16	2 ^d			3.7	1.55	0.82

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
NGC 5170	13 29	-17 57	3 ^e			5.0	1.92	0.91
UGC 8498	13 30	+31 37	4			3.0	1.44	0.46
NGC 5188	13 31	-34 47	5.1			2.6	1.48	0.43
ESO 21- 4	13 32	-77 50	3 ^d			-0.4	1.35	0.35
NGC 5229	13 34	+47 54	4			6.7	1.52	0.77
NGC 5221	13 34	+13 49	5.1			3.0	1.38	0.48
ESO 40- 12	13 35	-77 38	2			3.6	1.36	0.76
ESO 509- 74	13 35	-24 04	4			3.0 ⁵	1.37	0.72
NGC 5220	13 35	-33 27	4 ^d		SA II	1.0	1.37	0.53
IC 902	13 36	+49 57	4			3.0	1.34	0.72
ESO 383- 48	13 38	-33 31	2 ^d		SDB	-2.0	1.32	0.80
ESO 324- 44	13 38	-39 50	5.1			6.0	1.37	0.39
IC 4310	13 38	-25 50	3			-2.0	1.34	0.44
NGC 5290	13 45	+41 42	4		SA II	3.7	1.55	0.62
NGC 5297	13 46	+43 52	5.1			4.5	1.75	0.65
NGC 5301	13 46	+46 06	5.4			4.0	1.62	0.69
NGC 5308	13 47	+60 58	4			-3.0	1.57	0.74
UGC 8726	13 47	+40 29	5.4			7.0	1.35	0.72
UGC 8737	13 48	+68 05	2	3		4.0	1.35	0.72
ESO 577- 38	13 48	-18 52	5.5			7.0	1.32	1.27
MCG -1-35- 13	13 48	-07 11	2+			3.0	1.33	0.68
ESO 383- 91	13 50	-37 17	4			7.0	1.38	0.98
NGC 5346	13 53	+39 34	5.1			6.0	1.30	0.38
NGC 5353	13 53	+40 17	5.1			-2.0	1.34	0.30
NGC 5348	13 54	+05 13	4			3.5	1.55	0.86
NGC 5362	13 54	+41 18	5.1			3.0	1.36	0.35
NGC 5356	13 54	+05 20	3			3.5	1.49	0.56
NGC 5379	13 55	+59 44	5.1			-2.0	1.34	0.36
NGC 5389	13 56	+59 44	2		S	0.0	1.54	0.56
NGC 5365A	13 56	-44 00	3			3.0	1.41	0.71
IC 4351	13 57	-29 18	4 ^d		SA II	3.0	1.78	0.81
NGC 5403	13 59	+38 10	3			3.0	1.49	0.56
ESO 221- 22	14 00	-48 16	5.2			7.0	1.42	0.83
NGC 5422	14 00	+55 09	4			-2.0	1.59	0.72
NGC 5443	14 02	+55 48	5.1			3.0	1.43	0.43
NGC 5448	14 02	+49 10	5.1			1.0	1.60	0.35
NGC 5440	14 03	+34 45	5.1			1.0	1.49	0.41
NGC 5475	14 05	+55 44	4 ^a		SA III	0.5	1.31	0.61
NGC 5470	14 06	+06 01	3	4		3.0	1.40	0.83
NGC 5488	14 08	-33 18	4			3.5	1.53	0.53
ESO 446- 18	14 08	-29 34	1 ^d			3.0	1.37	1.01
UGC 9071	14 10	+54 12	4			6.0	1.32	0.72
UGC 9057	14 10	-02 34	5.1			7.3	1.48	0.48
NGC 5492	14 10	+19 36	4	4		3.0	1.38	0.55
NGC 5496	14 11	-01 09	5.2			7.0	1.67	0.72
UGC 9088	14 11	+38 11	4			0.0	1.41	0.58
ESO 97- 13	14 13	-65 20	5.1			3.0	1.84	0.36
NGC 5506	14 13	-03 12	1 ^b			1.0	1.45	0.52
ESO 271- 22	14 13	-45 24	5.2			5.8	1.38	0.81
UGC 9113	14 14	+35 25	3			5 ¹	1.32	0.65
NGC 5523	14 14	+25 19	3			6.0	1.66	0.56
NGC 5529	14 15	+36 13	1		SA II, S	5.0	1.79	0.87
IC 4377	14 16	-75 38	4			-1.0	1.43	0.48
IC 4393	14 17	-31 20	4 ^d			6.0	1.38	0.97
NGC 5560	14 20	+03 59	4			3.0	1.57	0.73
NGC 5566	14 20	+03 55	5.1			2.0	1.82	0.48

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
UGC 9182	14 20	+21 56	3			7.0	1.39	0.63
NGC 5577	14 21	+03 26	4			4.0	1.53	0.53
IC 4402	14 21	-46 17	5.2			3.0	1.62	0.69
NGC 5587	14 22	+13 55	5.1			0.0	1.42	0.50
NGC 5616	14 24	+36 27	4			4.0	1.33	0.69
NGC 5610	14 24	+24 36	5.1			2.0	1.31	0.48
UGC 9242	14 25	+39 32	4			7.0	1.70	1.22
UGC 9249	14 26	+08 40	4			7.0	1.32	0.86
ESO 1- 6	14 27	-87 46	5.2			5.6	1.48	0.63
NGC 5673	14 31	+49 57	3			5.3	1.39	0.59
IC 1029	14 32	+49 54	4			3.0	1.45	0.71
NGC 5675	14 32	+36 18	5.1			0 ²	1.45	0.46
IC 4453	14 34	-27 31	4			-2.0	1.35	0.37
NGC 5689	14 35	+48 44	3			0.0	1.54	0.56
UGC 9389	14 35	+12 54	4			3.0	1.36	0.51
NGC 5670	14 35	-45 57	5.1			-2.0	1.35	0.37
UGC 9394	14 35	+13 09	5.1			6.0	1.35	0.50
NGC 5707	14 37	+51 33	4	4	SA II	2.0	1.41	0.70
NGC 5690	14 37	+02 17	3			5.0	1.53	0.53
NGC 5714	14 38	+46 38	4			6.0	1.51	0.91
IC 4468	14 38	-22 22	5.1			4.8	1.34	0.53
UGC 9448	14 38	+51 09	3			3.0	1.34	0.95
IC 4472	14 40	-44 18	3			5.0	1.33	0.65
ESO 512- 12	14 40	-25 46	1 ^d		SDB	3.0	1.44	0.96
NGC 5719	14 40	-00 19	3			2.0	1.51	0.44
NGC 5729	14 42	-09 00	5.2			3.0	1.41	0.59
IC 1048	14 42	+04 53	4			3 ¹	1.35	0.49
ESO 222- 15	14 44	-49 24	5.6			6.0	1.32	0.48
NGC 5746	14 44	+01 57	1		J P, SA I, S	3.0	1.87	0.75
ESO 447- 36	14 46	-27 42	1 ^b			3.7	1.30	0.85
MCG -3-38- 8	14 47	-17 26	4			5.6	1.50	0.78
IC 1055	14 47	-13 43	5.1			3.0	1.30	0.46
ESO 580- 29	14 47	-19 45	3			5.3	1.33	0.79
NGC 5756	14 47	-14 51	5.1			3.5	1.32	0.35
IC 4484	14 47	-73 18	5.2			5.7	1.41	0.90
UGC 9537	14 48	+34 59	4			3.0	1.37	0.65
NGC 5777	14 51	+58 58	4			3.0	1.49	0.86
NGC 5775	14 53	+03 32	2 ^e	3		5.0	1.62	0.62
ESO 327- 31	14 55	-38 16	5.2			5.2	1.31	0.72
ESO 581- 4	14 57	-18 27	3			6.3	1.32	0.79
ESO 386- 43	14 58	-37 33	5.1			3.5	1.40	0.46
NGC 5792	14 58	-01 05	5.1			3.0	1.84	0.60
ESO 327- 39	14 59	-42 34	4			3.0	1.32	0.65
MCG -2-38- 27	15 01	-15 7	4			7.0	1.39	0.68
ESO 328- 5	15 01	-37 59	4			0.8	1.30	0.73
NGC 5838	15 05	+02 6	4			-3.0	1.62	0.45
NGC 5866	15 06	+55 45	4			-1.0	1.67	0.38
NGC 5859	15 07	+19 34	2	2	Sh	4.0	1.46	0.50
NGC 5854	15 07	+02 34	3			-1.0	1.44	0.54
ESO 223- 12	15 09	-52 33	5.6			4.1	1.37	0.69
NGC 5864	15 09	+03 3	2	3	SA I	-2.0	1.44	0.49
NGC 5879	15 09	+57 0	4			4.0	1.62	0.49
UGC 9759	15 10	+55 20	2	2		-0.5 ¹	1.30	0.67
IC 4522	15 11	-75 51	3			3.0	1.46	0.49
UGC 9760	15 12	+01 41	5.2			7.0	1.39	1.03
ESO 581- 25	15 13	-20 40	3			6.6	1.55	0.70

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
NGC 5878	15 13	−14 16	5.1			3.0	1.54	0.38
UGC 9780	15 14	+44 35	2			6.0	1.32	0.91
ESO 274- 1	15 14	−46 48	5.2			6.7	2.04	0.84
NGC 5899	15 15	+42 2	5.1			5.0	1.50	0.43
NGC 5907	15 15	+56 19	4			5.0	2.10	0.96
NGC 5908	15 16	+55 24	4 ^a	4	SA II	3.0	1.51	0.42
ESO 328- 41	15 18	−38 30	5.2			2.6	1.40	0.68
ESO 514- 5	15 19	−23 49	2 ^b			1.0	1.32	0.52
NGC 5916	15 21	−13 10	5.3			1.0	1.45	0.48
UGC 9829	15 23	−01 20	5.1			3.0	1.34	0.38
UGC 9841	15 25	+18 16	2	2		4.0	1.39	0.77
UGC 9853	15 25	+52 26	3	4		3.0	1.36	0.76
UGC 9856	15 26	+41 17	2			7.0	1.34	1.00
UGC 9858	15 26	+40 33	4	4		4.0	1.63	0.71
IC 4541	15 29	−70 35	4			3.0	1.37	0.62
NGC 5951	15 33	+15 0	2			5.3	1.54	0.62
NGC 5965	15 34	+56 41	1		SA II, S	3.0	1.72	0.85
NGC 5981	15 37	+59 23	4	4		5.0	1.45	0.79
NGC 5987	15 39	+58 4	4			3.0	1.62	0.50
UGC 9977	15 41	+00 42	3 ^d			5.3	1.58	0.91
NGC 5984	15 42	+14 13	2			7.0	1.46	0.58
ESO 99- 8	15 45	−66 17	5.2			7.0	1.36	1.06
NGC 6011	15 46	+72 10	5.1			3.0	1.30	0.44
UGC 10043	15 48	+21 52	4			4.0	1.38	0.82
NGC 6015	15 51	+62 18	5.1			6.0	1.73	0.40
ESO 515- 13	15 53	−27 17	4			3.0	1.30	0.48
IC 1151	15 58	+17 26	5.1			5.0	1.39	0.47
IC 4585	16 00	−66 19	5.1			3.0	1.33	0.52
ESO 136- 16	16 03	−60 58	5.6			5.0	1.44	0.80
UGC 10169	16 04	+14 49	5.3			−2.0	1.32	0.74
UGC 10214	16 06	+55 25	5.1			5.0	1.56	0.64
IC 1197	16 08	+07 32	4			6.0	1.46	0.78
UGC 10227	16 08	+36 36	4	3		6.0	1.36	0.97
ESO 137- 3	16 13	−61 00	5.6			5.5 ⁵	1.54	0.43
UGC 10288	16 14	−00 12	2 ^e			5.3	1.68	0.92
UGC 10297	16 15	+18 54	5.4			5.0	1.32	0.93
UGC 10306	16 16	+00 14	5.1			3.0	1.32	0.44
IC 4595	16 20	−70 08	5.2			5.0	1.43	0.70
ESO 137- 18	16 20	−60 29	5.6			5.0	1.50	0.45
NGC 6118	16 21	−02 16	5.1			6.0	1.67	0.37
UGC 10447	16 28	+80 21	3	4		3.0	1.30	0.62
UGC 10413	16 29	+21 20	5.1			6.0	1.35	0.47
ESO 452- 7	16 32	−28 05	5.6			2.5	1.42	0.41
NGC 6181	16 32	+19 49	5.1			5.0	1.40	0.35
ESO 137- 37	16 39	−59 53	5.6			5 ⁶	1.32	0.94
ESO 137- 38	16 40	−60 23	5.6			4.2	1.42	0.44
NGC 6207	16 43	+36 49	5.1			5.0	1.47	0.36
UGC 10536	16 44	+62 04	5.1			3.0	1.34	0.37
NGC 6248	16 46	+70 21	5.1			7.0	1.50	0.43
UGC 10561	16 46	+62 48	4			3.0	1.38	0.95
NGC 6239	16 50	+42 44	5.1			3.0	1.41	0.35
NGC 6255	16 54	+36 30	5.1			6.0	1.56	0.37
UGC 10610**	16 54	+43 03	5.3			?	1.34	0.38
ESO 138- 7	16 56	−62 24	5.1			4.0	1.33	0.39
ESO 138- 14	17 06	−62 05	5.2			6.5	1.62	0.90
NGC 6310	17 07	+60 59	2	2		3.0	1.30	0.67

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
IC 4635	17 15	−77 29	2			3.3	1.48	0.64
NGC 6361	17 18	+60 36	3	4		3.0	1.34	0.54
ESO 138- 24	17 24	−59 22	5.2			5.0	1.38	0.63
NGC 6395	17 26	+71 05	5.1			6.0	1.38	0.51
NGC 6368	17 27	+11 32	4			3.0	1.58	0.59
IC 1265	17 36	+42 05	5.1			2.0	1.31	0.37
NGC 6434	17 36	+72 05	5.1			4.0	1.37	0.38
IC 4656	17 37	−63 43	5.2			4.5	1.36	0.59
NGC 6433	17 43	+36 47	4			3.0	1.30	0.58
ESO 139- 21	17 44	−60 58	3			3.0	1.30	0.66
UGC 10972	17 46	+26 32	4			6.0	1.40	0.57
NGC 6503	17 49	+70 08	5.1			6.0	1.85	0.47
NGC 6504	17 56	+33 12	4		SA I	2 ¹	1.35	0.68
UGC 11057	17 57	+12 10	5.1			6.0	1.30	0.39
ESO 139- 49	18 00	−59 08	5.2			6.0	1.36	0.54
UGC 11093	18 01	+06 58	5.2			6.0	1.50	0.78
UGC 11132	18 09	+38 47	4			3.0	1.32	0.95
UGC 11140	18 11	+39 10	4			6.0	1.38	0.76
IC 1279	18 11	+36 00	4			3.0	1.41	0.66
IC 4679	18 11	−56 15	5.1			6.0	1.34	0.40
NGC 6621	18 13	+68 21	5.1			3.0	1.32	0.41
IC 4694	18 15	−58 12	5.1			5.0	1.35	0.43
IC 4696	18 20	−64 44	5.1			4.2	1.37	0.41
NGC 6636	18 22	+66 36	5.3			4 ²	1.34	0.78
ESO 395- 2	18 22	−35 40	5.6			3.0	1.31	0.80
ESO 280- 13	18 23	−43 08	5.6			4.0	1.46	0.65
UGC 11230	18 23	+65 19	4			7.0	1.45	1.18
IC 4720	18 33	−58 24	5.1			6.3	1.40	0.46
IC 4721	18 34	−58 29	5.2			6.0	1.72	0.55
NGC 6690	18 34	+70 31	5.1			7.0	1.58	0.48
UGC 11291	18 36	+30 49	5.2			7.0	1.35	0.64
IC 4728	18 37	−62 31	5.1			2.0	1.30	0.44
NGC 6654A	18 39	+73 34	5.2			7.2	1.42	0.54
UGC 11325	18 40	+36 09	4			3.0	1.36	0.68
IC 4745	18 42	−64 56	2 ^c		SDB	2.0	1.33	0.39
ESO 336- 16	18 47	−41 43	5.6			5.0	1.31	0.40
ESO 396- 7	18 54	−34 36	5.6			5.0	1.30	0.45
NGC 6707	18 55	−53 49	5.1			5.0	1.32	0.35
IC 4806	19 01	−57 31	4			3.0	1.37	0.61
NGC 6725	19 01	−53 51	2			−2.0	1.34	0.66
IC 4810	19 02	−56 09	2			6.8	1.54	0.96
UGC 11394	19 03	+27 36	4			6.0	1.31	0.93
NGC 6722	19 03	−64 53	1 ^d		J P, SDB	3.0	1.46	0.84
IC 4819	19 07	−59 27	2			6.0	1.46	0.95
NGC 6739	19 07	−61 22	4			−2.2	1.38	0.45
UGC 11411A	19 10	+60 07	5.5			6.0	1.32	0.85
IC 4827	19 13	−60 51	3			2.3	1.45	0.76
IC 4832	19 14	−56 36	3			1.0	1.35	0.68
IC 4831	19 14	−62 16	4			2.3	1.53	0.58
IC 4837A	19 15	−54 07	4			2.5	1.61	0.81
ESO 184- 51	19 16	−53 42	4			4.0	1.32	0.49
NGC 6771	19 18	−60 32	1 ^b	1	J P, SDB	−1.0	1.36	0.67
ESO 184- 63	19 23	−55 03	4			3.0	1.31	0.65
UGC 11435	19 23	+55 59	4			6.0	1.30	0.76
NGC 6788	19 26	−54 57	4			2.0	1.46	0.52
UGC 11455	19 29	+72 06	4	4		6.0	1.42	0.87

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
ESO 142- 19	19 33	-58 06	4 ^d			3.0	1.61	0.78
IC 4872	19 35	-57 31	3 ^d			6.7	1.54	0.98
ESO 460- 18	19 38	-29 48	5.1			4.0	1.31	0.46
MCG -3-50- 3	19 42	-15 52	5.2			5.0	1.31	0.79
NGC 6810	19 43	-58 39	4 ^d			2.0	1.50	0.55
NGC 6816	19 43	-28 29	5.1			4.0	1.35	0.36
ESO 460- 31	19 44	-27 24	4 ^d			4.5	1.47	1.03
NGC 6835	19 54	-12 34	4			1.0	1.37	0.66
UGC 11498	19 57	+05 53	5.2			3.0	1.50	0.51
NGC 6848	20 02	-56 05	3		SA III	0.5	1.39	0.38
UGC 11510	20 02	+53 53	5.1			7.0	1.31	0.51
NGC 6850	20 03	-54 50	5.1			-1.0	1.33	0.30
ESO 105- 22	20 03	-65 04	5.5			7 ²	1.30	1.19
NGC 6861D	20 08	-48 12	4			-3.0	1.37	0.48
NGC 6867	20 10	-54 47	5.1			4.0	1.30	0.47
NGC 6875A	20 11	-46 08	3			4.0	1.43	0.77
NGC 6872	20 16	-70 46	5.1			3.0	1.78	0.54
ESO 340- 8	20 17	-40 55	4			6.0	1.35	1.38
NGC 6887	20 17	-52 47	5.1			4.0	1.53	0.42
ESO 340- 9	20 17	-38 40	5.5			7.0	1.35	0.98
UGC 11537	20 18	-00 09	5.2			4.7	1.38	0.55
NGC 6880	20 19	-70 51	2			-1.0	1.31	0.36
IC 4992	20 23	-71 33	3			5.2	1.35	0.99
IC 4946	20 23	-43 59	5.1			-0.1	1.39	0.38
ESO 340- 26	20 24	-41 30	2			5.0	1.31	0.94
ESO 400- 20	20 25	-34 11	5.2			3.0	1.33	0.51
UGC 11568	20 28	+10 45	5.2			6.0	1.33	0.54
UGC 11571	20 29	+10 40	5.2			6.0	1.31	0.56
NGC 6928	20 32	+09 55	4			2.0	1.30	0.49
NGC 6925	20 34	-31 58	4			4.0	1.65	0.56
MCG -1-52- 8	20 35	-06 14	5.2			5.0	1.31	0.87
MCG -1-52- 16	20 38	-05 38	4			1.0	1.39	0.49
IC 5039	20 43	-29 51	5.1			4.3	1.38	0.60
UGC 11635	20 43	+80 09	5.1			4.0	1.46	0.37
NGC 6948	20 43	-53 21	4 ^d			1.4	1.34	0.36
ESO 285- 48	20 44	-45 58	5.1			6.0	1.39	0.39
ESO 234- 69	20 45	-51 23	5.1			5.0	1.32	0.49
IC 5026	20 48	-78 04	4			7.0	1.37	0.94
MCG -1-53- 12	20 49	-07 01	4			5.0	1.48	0.92
IC 5052	20 52	-69 12	5.4 ^d			7.0	1.77	0.88
IC 5054	20 53	-71 01	4			1.0	1.30	0.72
ESO 286- 18	20 57	-43 22	3			4.0	1.39	0.83
IC 5071	21 01	-72 38	4			5.0	1.53	0.61
IC 5078	21 02	-16 48	4			5.0	1.61	0.56
NGC 7013	21 03	+29 53	4			0.0	1.60	0.45
ESO 235- 53	21 05	-47 47	2			3.0	1.39	0.73
ESO 235- 57	21 06	-48 10	4			4.0	1.42	0.58
ESO 402- 10	21 08	-33 14	5.1			-2.0	1.30	0.37
NGC 7020	21 11	-64 01	5.1			-1.0	1.57	0.31
ESO 187- 58	21 12	-57 16	2			4.7	1.32	0.84
ESO 107- 15	21 13	-63 20	3			6.0	1.36	0.76
NGC 7041	21 16	-48 21	4		SA III	-2.7	1.56	0.39
IC 5096	21 18	-63 45	4 ^d			4.0	1.50	0.80
ESO 145- 4	21 18	-57 38	4			5.4	1.35	0.64
ESO 402- 26	21 22	-36 40	5.1			2.4	1.44	0.54
ESO 287- 13	21 23	-45 46	4			4.0	1.50	0.56

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
NGC 7064	21 29	-52 45	5.4 ^d			5.0	1.58	0.79
ESO 48- 2	21 36	-76 20	2			3.0	1.37	0.60
NGC 7090	21 36	-54 33	3			5.0	1.87	0.77
MCG 2-55- 7	21 40	+12 21	4			3 ⁶	1.37	0.63
ESO 531- 22	21 40	-26 31	3			4.0	1.41	1.00
NGC 7124	21 48	-50 33	5.1			4.0	1.47	0.37
IC 5139	21 50	-30 59	5.1			-3.0	1.32	0.31
UGC 11838	21 52	+28 18	5.5			7.0	1.31	1.03
IC 5140	21 54	-67 19	2			6.5	1.30	0.86
NGC 7151	21 55	-50 39	5.1			5.8	1.47	0.40
UGC 11859	21 58	+01 00	4 ^d			4.0	1.49	1.13
ESO 288- 25	21 59	-43 52	4			4.0	1.36	0.87
NGC 7162	21 59	-43 18	5.1			5.0	1.45	0.43
NGC 7166	22 00	-43 23	4			-3.0	1.39	0.43
ESO 404- 18	22 01	-32 34	4			7.0	1.37	1.13
NGC 7183	22 02	-18 54	3			-0.5	1.58	0.52
NGC 7184	22 02	-20 48	3		Sh	5.0	1.78	0.61
IC 5156	22 03	-33 50	5.1			1.6	1.35	0.44
ESO 404- 27	22 03	-32 17	5.1			4.7	1.35	0.39
UGC 11893	22 04	+35 56	3	4		7.0	1.31	0.81
NGC 7179	22 04	-64 02	5.1			3.7	1.31	0.40
NGC 7218	22 10	-16 39	5.1			6.0	1.40	0.35
IC 5171	22 10	-46 04	3			4.0	1.48	0.83
ESO 146- 14	22 13	-62 04	4			7.0	1.45	1.22
IC 5181	22 13	-46 01	4		SA II	-2.0	1.42	0.51
NGC 7232A	22 13	-45 53	3			2.0	1.34	0.73
ESO 533- 4	22 14	-26 56	3 ^d			4.8	1.37	0.86
IC 5176	22 14	-66 50	4 ^d		SA III	3.9	1.65	0.92
NGC 7232	22 15	-45 51	4			1.0	1.41	0.48
NGC 7241	22 15	+19 13	5.2			4.0	1.53	0.49
ESO 237- 49	22 16	-47 39	3			5.8	1.33	0.80
ESO 289- 10	22 16	-47 07	4			6.7	1.30	0.77
UGC 11973	22 16	+41 30	2	2		4.0	1.51	0.58
UGC 11994	22 20	+33 17	4			4.0	1.35	0.92
NGC 7264	22 22	+36 23	4	4		3.0	1.35	0.84
ESO 27- 8	22 23	-79 59	5.1			5.0	1.43	0.40
ESO 467- 51	22 23	-28 58	3			6.0	1.45	1.04
NGC 7282	22 25	+40 18	5.1			3.0	1.39	0.38
UGC 12073	22 32	+39 12	5.1			3.0	1.32	0.41
NGC 7307	22 33	-40 56	5.4			5.6	1.55	0.60
NGC 7331	22 37	+34 25	5.1			3.0	2.02	0.45
NGC 7332	22 37	+23 47	3 ^f	3	J P, SA II	-2.0	1.61	0.56
NGC 7328	22 37	+10 31	5.1			2.0	1.31	0.44
NGC 7339	22 37	+23 47	3 ^f	3		4.0	1.48	0.61
ESO 534- 9	22 38	-25 50	4			2.0	1.46	0.56
NGC 7341	22 39	-22 39	5.1			1.8	1.38	0.40
NGC 7361	22 42	-30 03	4			5.0	1.58	0.60
IC 5244	22 44	-64 02	3			3.0	1.46	0.83
NGC 7359	22 44	-23 41	4			-2.0	1.37	0.61
NGC 7368	22 45	-39 20	3			3.4	1.47	0.74
MCG -2-58- 3	22 46	-14 10	4			7.0	1.30	0.82
IC 5249	22 47	-64 49	4			7.0	1.56	0.96
UGC 12190	22 48	+28 17	4			5.0	1.32	0.96
UGC 12221	22 50	+82 52	5.1			7.0	1.37	0.54
NGC 7400	22 54	-45 20	4			4.0	1.42	0.70
NGC 7410	22 55	-39 39	3			1.0	1.72	0.51

Table 6 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
NGC 7416	22 55	−05 29	5.1			3.0	1.50	0.55
IC 5267A	22 55	−43 26	5.1			4.3	1.38	0.48
IC 5269B	22 56	−36 14	4			5.5	1.61	0.72
IC 5264	22 56	−36 33	4			2.3	1.39	0.69
MCG -2-58- 11	22 56	−08 58	4			5.0	1.30	0.68
IC 5270	22 57	−35 51	3			5.7	1.50	0.71
IC 5271	22 58	−33 44	3			3.0	1.42	0.49
IC 5269C	23 00	−35 22	5.1			6.9	1.33	0.38
UGC 12308	23 01	+14 20	4			6.0	1.41	0.67
ESO 290- 35	23 01	−46 38	4			4.7	1.30	0.80
NGC 7463	23 01	+15 58	5.1			3.0	1.46	0.63
NGC 7456	23 02	−39 34	5.1			6.0	1.74	0.53
NGC 7462	23 02	−40 50	3			4.0	1.62	0.76
UGC 12344	23 04	+18 43	5.5			7.0	1.33	0.75
ESO 469- 15	23 08	−30 51	4			3.0	1.31	0.95
NGC 7497	23 09	+18 10	5.1			6.5	1.69	0.63
ESO 407- 7	23 09	−36 25	4			3.0	1.45	0.84
MCG -2-59- 4	23 10	−08 41	5.1			5.0	1.30	0.91
NGC 2573A	23 12	−89 07	5.2			3.0	1.33	0.45
UGC 12423	23 13	+06 24	4 ^f			5.0	1.54	0.98
UGC 12430	23 13	+29 00	3			7.0	1.35	1.02
ESO 347- 2	23 14	−40 37	5.5			6.8	1.31	0.85
NGC 7537	23 14	+04 29	4			4.0	1.35	0.59
NGC 7541	23 14	+04 32	5.1			4.0	1.54	0.45
NGC 7531	23 14	−43 35	5.1			4.0	1.65	0.40
NGC 7549	23 15	+19 02	5.1			6.0	1.45	0.58
NGC 7582	23 18	−42 22	1			2.0	1.70	0.38
NGC 7600	23 18	−07 34	5.1			−3.0	1.40	0.36
NGC 7590	23 18	−42 14	5.1			4.0	1.43	0.42
NGC 7606	23 19	−08 29	5.1			3.0	1.73	0.40
NGC 7599	23 19	−42 15	5.1			5.0	1.64	0.51
UGC 12506	23 19	+16 04	4	4		6.0	1.46	1.17
NGC 7632	23 22	−42 28	5.1			−2.0	1.34	0.31
NGC 7640	23 22	+40 50	2	2		5.0	2.02	0.72
NGC 7690	23 33	−51 41	5.1			3.0	1.35	0.41
NGC 7700	23 34	−02 57	4			−1.0	1.31	0.66
NGC 7703	23 34	+16 04	4			−2.0	1.35	0.68
NGC 7709	23 35	−16 42	3			−2.0	1.36	0.55
NGC 7711	23 35	+15 18	5.1			−2.0	1.42	0.32
UGC 12693	23 35	+32 23	4			6.0	1.34	1.12
NGC 7713	23 36	−37 56	5.1			7.0	1.65	0.39
ESO 148- 20	23 36	−57 37	2			7.0	1.34	1.10
ESO 240- 10	23 37	−47 30	5.1			−2.0	1.60	0.33
ESO 240- 11	23 37	−47 43	4 ^d		SA II	4.8	1.72	1.02
NGC 7721	23 38	−06 31	5.1			5.0	1.55	0.39
ESO 471- 2	23 40	−28 20	5.1			4.2	1.32	0.45
ESO 292- 14	23 42	−44 54	4			6 ⁶	1.40	0.88
ESO 12- 4	23 44	−80 10	4			5.0	1.30	0.98
NGC 7771	23 51	+20 06	5.1	5.1		1.0	1.40	0.39

Notes to the table:

Col. (1): Principal name of the galaxy using following order: NGC, IC, ESO, UGC, MCG, UGCA, and PGC.

*: NGC 2788A = ESO 60- 24 (NGC number is not in RC3).

** : UGC 10610 = MCG 7-35- 4 (UGC number is not in RC3).

Col. (2): RA in hours and minutes (leaving the seconds).

Col. (3): DEC in degree and minutes (leaving the seconds).

Col. (4): Bulge type as defined in section 3. Numbers marked with a letter are confirmed by CCD observations.

^a: Calar Alto, 1.2m, 1996

^b: ESO/La Silla, 0.9m, 1998

^c: ESO/La Silla, 1.54m, 1996, 1998

^d: ESO/La Silla, 2.2m, 1985, 1987, 1990

^e: ESO/La Silla, NTT, 1991

^f: Lowell, 1.06m, 1989, 1990

^g: ESO/ST-ECF Archive, VLT, 1999

Col. (5): Bulge type in the NIR (for details of the observations cf. Paper II).

Col. (6): Earlier detections and classifications:

J P: Jarvis (1986), peanut-shaped

J B: Jarvis (1986), box-shaped

SA I: SA87 (1987), class I

SA II: SA87 (1987), class II

SA III: SA87 (1987), class III

S: Shaw (1987)

SDB: Shaw et al. (1990) (only literature sample)

Sh: Shaw (1993a) (only galaxies not listed in S)

Col. (7): Morphological type from RC3, except for

¹: Skiff (1999)

²: Own classification, only bulges of type 5 (checked by 12 galaxies classified by Skiff (1999), $T_{skiff} - T_{own} \leq |1|$).

³: NGC 3079: Morphological type (SBS5) differs from T (7.0) (both RC3).

⁴: UGC 8032: Morphological type from Skiff (1999) and Haynes et al. (1990). The T value of Haynes et al. is based on Kraan-Korteweg (1982) who refers to a priv. communication with Sandage (1980). Yasuda et al. (1995, 1997) gave a wrong classification $T = 5$. In 1995 they refer to VCC (Binggeli et al. 1985) and in 1997 to RC3, but in both catalogues the type of UGC 8032 is not listed.

⁵: “The Surface Photometry Catalogue of the ESO-Uppsala Galaxies” (Lauberts & Valentijn 1989).

⁶: NASA Extragalactic Database (NED).

⁷: “The ESO/Uppsala Survey of the ESO (B) Atlas” (Lauberts 1982).

Col. (8): Decimal logarithm of the 25 (B) isophotal major diameter from the RC3.

Col. (9): Decimal logarithm of the 25 (B) isophotal axis ratio (a/b) from the RC3.

Table 7. List of galaxies with bulge classification not included in our selected RC3-sample

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) $\log D_{25}$	(9) $\log R_{25}$
ESO 112- 4	00 28	-58 06	2 ^d		SDB	5.6	1.12	0.85
ESO 150- 14	00 36	-56 54	4 ^d			-0.7	1.28	0.99
ESO 195- 5	00 50	-52 07	4		J B	1 ⁶	1.04 ⁶	0.74 ⁶
ESO 151- 4	00 56	-53 11	1		J P, SDB	-2.0	1.13	0.56
ESO 113- 6	01 01	-59 44	3 ^d			5.0	1.22	0.97
NGC 489	01 21	+09 12	4		SA II	-2 ⁶	1.22	0.65
ESO 244- 48	01 39	-47 07	4 ^d			3.0	1.14	0.82
ESO 244- 49	01 39	-46 59	3		J P	-0.6	1.21	0.94
NGC 1030	02 39	+18 01	3 ^f			-1 ¹	1.20	0.38
NGC 1175	03 04	+42 20	2+	2+	J B	-1.0	1.29	0.50
ESO 205- 9	05 57	-52 22	4		J B	3.7	1.27	0.86
NGC 2191	06 08	-52 30	5.1			-2.0	1.24	0.31
NGC 2769	09 10	+50 25	2 ^f			1.0	1.24	0.62
ESO 433- 19	09 24	-28 10	1		J B, SDB	3 ⁶	1.23 ⁶	0.75 ⁶
IC 2560	10 16	-33 33	2 ^d			3.3	1.50	0.20
ESO 376- 9	10 42	-33 14	4 ^d			-2.0	1.19	0.56
ESO 318- 2	10 42	-40 34	4 ^d			1.0	1.14	0.61
ESO 376- 23	10 51	-35 28	4 ^d			5.0	1.25	0.95
NGC 3524	11 06	+11 23	4		SA II	0.0	1.21	0.54
NGC 3564	11 10	-37 33	4			-2.0	1.25	0.37
ESO 377- 29	11 14	-33 54	4			-2.0	1.18	0.40
NGC 3627	11 20	+12 59	5.1		Sh	3.0	1.96	0.34
NGC 3650	11 22	+20 42	2		SDB	3.0	1.23	0.75
ESO 319- 26	11 30	-41 03	4 ^d			5.3	1.17	1.02
NGC 3762	11 37	+61 45	3 ^f		SA II, SDB	1.0	1.27	0.58
NGC 3838	11 44	+57 57	5.1		SA III	-0.4	1.19	0.39
ESO 378- 20	11 47	-37 33	5.1			-2.1	1.11	0.33
ESO 572- 44	12 01	-20 29	1 ^d		J P, SDB	3.0	1.22	0.77
IC 760	12 05	-29 17	3	3	J P	-1.7	1.22	0.53
ESO 321- 10	12 11	-38 32	1 ^d		SDB	0.0	1.27	0.93
NGC 4215	12 15	+06 24	3			-1.0	1.27	0.42
ESO 506- 3	12 21	-25 04	3 ^d			2.3	1.15	0.48
ESO 506- 33	12 40	-25 19	4 ^d			-2.0	1.26	0.43
ESO 322- 51	12 40	-41 36	3			-3.0	1.23	0.54
NGC 4645B	12 43	-41 21	4			-2.2	1.28	0.42
ESO 322-100	12 49	-41 27	2 ^d			-2.0	0.97	0.37
ESO 575- 59	13 07	-19 23	4 ^d			-0.8	1.27	0.66
NGC 5014	13 11	+36 16	4	4	SA II	1.0	1.23	0.46
NGC 5193A	13 31	-33 14	5.2 ^d		SDB	-1.9	1.09	0.52
NGC 5253	13 39	-31 38	4.0		SA III	10.0	1.70	0.41
ESO 445- 49	13 49	-31 09	4 ^d			0.0	1.15	0.41
ESO 383- 85	13 49	-33 05	3 ^d			0 ⁶	1.04 ⁶	0.74 ⁶
ESO 383- 86	13 49	-33 11	3 ^d			-2 ⁶	1.00 ⁶	0.70 ⁶
ESO 445- 63	13 52	-30 49	1 ^b		J P	5 ⁵	1.08 ⁶	0.60 ⁶
NGC 5333	13 54	-48 30	5.1			-2.0	1.28	0.30
ESO 510- 13	13 55	-26 46	3 ^g			1.0	1.29	0.17
ESO 578- 25	14 08	-20 00	2 ^d			1 ⁶	1.20 ⁶	0.60 ⁶
ESO 510- 74	14 08	-26 56	3 ^d			1.3	1.23	0.67
ESO 385- 8	14 19	-34 51	4 ^d			7 ⁶	1.28 ⁶	0.98 ⁶
IC 4464	14 37	-37 36	3			-1.8	1.21	0.41
ESO 581- 6	14 58	-19 23	4 ^d			7.0	1.23	0.86
ESO 583- 8	15 57	-22 29	4 ^d		J B	6.0	1.18	0.97 ⁶
UGC 10205	16 06	+30 06	3	3		1.0	1.16	0.23
UGC 10535	16 46	+06 28	4 ^d			2.0	1.04	0.76
ESO 71- 1	17 59	-67 56	4 ^d			3 ⁵	0.99 ⁵	0.70 ⁵

Table 7 continued.

(1) Object	(2) RA (2000)	(3) DEC (2000)	(4) bulge type	(5) NIR bulge type	(6) earlier detections	(7) T	(8) log D_{25}	(9) log R_{25}
IC 4731	18 38	−62 56	5.1			−1.3	1.17	0.31
IC 4757	18 43	−57 09	2 ^b			−0.3	1.14	0.39
ESO 103- 59	18 45	−63 31	4		J B	−2 ⁶	1.14 ⁶	0.85 ⁶
IC 4767	18 47	−63 24	1 ^b		J P, SDB	−1.0	1.17	0.51
IC 4808	19 01	−45 18	5.1			5.0	1.29	0.37
NGC 6737	not a galaxy				SA I			
ESO 232- 21	19 44	−51 36	4			−1.5	1.24	0.74
ESO 398- 29	19 45	−34 44	1 ^b		J P	4 ⁶	1.23 ⁶	0.89 ⁶
ESO 105- 12	19 46	−65 14	4		J P	−2 ⁶	1.08 ⁶	0.78 ⁶
ESO 461- 6	19 51	−31 58	3 ^d			5.0	1.21	1.13
ESO 339- 16	20 00	−40 43	2 ^d		J P	1 ⁶	1.15 ⁶	0.85 ⁶
ESO 185- 53	20 03	−55 56	2		J P, SDB	−2.0	1.08	0.42
IC 4937	20 05	−56 15	1 ^d		J P, SDB	3.0	1.29	0.64
ESO 186- 36	20 21	−53 45	2			1.0	1.27	0.55
ESO 528- 17	20 33	−27 05	4 ^d			5.7	1.20	1.04
ESO 187- 8	20 43	−56 12	5.5 ^d			6.0	1.18	0.88
ESO 597- 36	20 48	−19 50	1 ^b		J P, SDB	−1.7	1.20	0.68
IC 5053	20 53	−71 08	4 ^d			1.0	1.20	0.43
ESO 235- 35	21 01	−52 00	2		J B	3.0	1.10	0.76
ESO 466- 1	21 42	−29 22	4 ^d			2.0	1.14	0.66
NGC 7123	21 50	−70 20	4 ^d		SA II	1.0	1.48	0.32
ESO 189- 12	21 55	−54 52	3 ^d		SDB	5.0	1.22	0.96
ESO 466- 46	22 02	−31 59	5.1		J P	−2 ⁵	1.08	0.50
IC 5199	22 19	−37 32	3 ^d		J B, SDB	3.0	1.19	0.85
IC 5267B	22 56	−43 45	2 ^d			−2.0	1.27	0.35
IC 5269	22 57	−36 01	5.1		SDB	−1.8	1.25	0.34
NGC 7443	23 00	−12 48	4			−1.0	1.17	0.42
ESO 604- 6	23 14	−20 59	3 ^d		SDB	4.0	1.27	0.85

Notes to the table:

Col. (1) - (7): s. Table 6

Col. (8): Decimal logarithm of the 25 (B) isophotal major axis diameter from RC3, except ⁵: ESO-IV, ⁶: NED.Col. (9): Decimal logarithm of the 25 (B) isophotal axis ratio (a/b) from RC3, except ⁵: ESO-IV, ⁶: NED.